

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07K 13/00, C12N 15/62	A1	(11) International Publication Number: WO 93/10151 (43) International Publication Date: 27 May 1993 (27.05.93)
<p>(21) International Application Number: PCT/US92/10272</p> <p>(22) International Filing Date: 20 November 1992 (20.11.92)</p> <p>(30) Priority data: 07/797,556 22 November 1991 (22.11.91) US</p> <p>(71) Applicant: IMMUNEX CORPORATION [US/US]; 51 University Street, Seattle, WA 98101 (US).</p> <p>(72) Inventor: GEARING, David, P. ; 4007 - 37th Avenue SW, Seattle, WA 98126 (US).</p> <p>(74) Agent: SEESSE, Kathryn, A.; Immunex Corporation, 51 University Street, Seattle, WA 98101 (US).</p>		<p>(81) Designated States: AU, CA, FI, JP, KR, NO, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE).</p> <p>Published <i>With international search report.</i></p>
<p>(54) Title: RECEPTOR FOR ONCOSTATIN M AND LEUKEMIA INHIBITORY FACTOR</p> <p>(57) Abstract</p> <p>A receptor protein comprising a gp130 polypeptide linked to a single-chain leukemia inhibitory factor receptor (LIF-R) polypeptide is capable of binding both oncostatin M and leukemia inhibitory factor (LIF). The receptor protein binds LIF with greater affinity than does the single-chain LIF-R polypeptide alone. The receptor may be produced as a fusion protein in recombinant cells. The gp130 polypeptide binds oncostatin M, but with lower affinity than does the inventive receptor protein.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Cote d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CN	China	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TC	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

WO 93/10151

PCT/US92/10272

1

TITLE

Receptor for Oncostatin M and Leukemia Inhibitory Factor

BACKGROUND OF THE INVENTION

Receptors that bind specific molecules (e.g., a hormone, drug, cytokine, or biochemical) have been identified on a multitude of cell types. Receptors are found on the cell surface or, in the case of soluble receptors, are released into the serum. Effort has been directed toward isolation and characterization of a number of receptors in order to study their physiological roles and to explore possible therapeutic uses. The binding of a particular target molecule by a soluble receptor administered to a patient may alleviate disorders mediated by the target molecule.

Certain receptors have been found to comprise two separate polypeptide chains associated in the form of a complex. Such two-chain receptors often bind the target molecule with greater affinity than that exhibited by one of the chains alone.

Leukemia inhibitory factor (LIF) is a polypeptide hormone that plays a central role in the regulation of diverse adult and embryonic systems. LIF acts on a variety of cell types and has multiple biological activities. The diversity in biological activity is reflected in the various synonyms of LIF, which include hepatocyte stimulating factor III (Baumann and Wong, *J. Immunol.* 143:1163 [1989]); cholinergic nerve differentiation factor (Yamamori et al., *Science* 246: 1412 [1990]); melanoma-derived lipoprotein lipase inhibitor (Mori et al., *Biochem. Biophys. Res. Comm.* 160:1085 [1989]); human interleukin for DA cells (Moreau et al., *Nature* 336:690 [1988]); differentiation factor (Tomida et al., *J. Biol. Chem.* 259:10978 [1984]); differentiation inhibitory factor (Abe et al., *J. Biol. Chem.* 264: 8941 [1989]); differentiation inhibitory activity (Smith and Hooper, *Devel. Biol.*; 121:1 [1987]); and differentiation retarding factor (Koopman and Cotton, *Exp. Cell. Res.* 154:233 [1984]).

The cloning of a leukemia inhibitory factor receptor (LIF-R) has been reported by Gearing et al. in *EMBO J.* 10:2839 (1991). This recombinant single-chain LIF-R polypeptide binds LIF, but with lower affinity than the naturally occurring LIF

WO 93/10151

PCT/US92/10272

2

receptors found on certain normal cells. A receptor that binds LIF with higher affinity than that exhibited by this cloned single chain LIF-R would be desirable for certain applications.

5 Oncostatin M is a secreted single-chain polypeptide cytokine that regulates the growth of certain tumor-derived and normal cell lines. Oncostatin M is produced by activated lymphoid cells. A number of cell types have been found to bind the oncostatin M protein. See, for example, Linsley et al., *J. Biol. Chem.*, 264: 4282 (1989). However, the isolation and characterization of an oncostatin M receptor have not been reported.

10

SUMMARY OF THE INVENTION

The present invention provides a receptor that has the property of binding both oncostatin M and leukemia inhibitory factor (LIF). The receptor comprises gp130
15 linked (preferably covalently) to leukemia inhibitory factor receptor (LIF-R). The gp130 polypeptide may be covalently linked to the LIF-R polypeptide by any suitable means, such as via a cross-linking reagent or a polypeptide linker. In one embodiment of the invention, the receptor is a fusion protein produced by recombinant DNA technology. Disorders mediated by either oncostatin M or LIF may be treated by administering a therapeutically effective amount of the inventive receptor to a patient
20 afflicted with such a disorder.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graph presenting the results of an LIF binding assay. Host cells
25 transfected with vector(s) encoding gp130 or LIF-R were assayed for the ability to bind LIF, as described in example 1.

Figure 2 is a graph presenting the results of an oncostatin M binding assay. Host cells transfected with vector(s) encoding gp130 or LIF-R were assayed for the ability to bind oncostatin M, as described in example 2.

Figure 3 is a graph depicting low affinity binding of oncostatin M to host cells
30 transfected with a gp130 encoding expression vector, as described in example 2.

Figure 4 schematically depicts a receptor of the present invention wherein Fc polypeptides derived from an antibody are used to link a gp130 fragment to an LIF-R fragment.

Figure 5 presents composite DNA and encoded amino acid sequences of a full
35 length LIF-R, determined by comparing the sequences of cDNA and genomic clones. The signal peptidase cleavage site is marked with a vertical arrow. The transmembrane region is heavily underlined. Potential N-linked glycosylation sites are marked with

WO 93/10151

PCT/US92/10272

3

asterisks. Hallmark residues associated with the hematopoietin family of receptors are shown boxed. The horizontal arrow marks the point at which genomic sequence was used to derive the 3' coding region of LIF-R, since the cDNA clones employed in determining this sequence terminated with a stretch of A nucleotides at this point.

5 Figure 6 presents the DNA and deduced amino acid sequences of cloned gp130 cDNA as reported by Hibi et al. in *Cell* 63:1149 (1990). A predicted signal sequence is underlined. The thick underline indicates a presumed transmembrane region. The sets of asterisks identify possible N-glycosylation sites.

10 Figure 7 presents Scatchard analyses that demonstrate the interaction of a soluble gp130/Fc fusion protein with soluble LIF-R/Fc in binding LIF and oncostatin M, as described in example 7.

DETAILED DESCRIPTION OF THE INVENTION

15 The present invention provides a receptor comprising gp130 covalently linked to leukemia inhibitory factor receptor (LIF-R). In another embodiment of the invention, the receptor comprises gp130 non-covalently complexed with LIF-R. The receptor is capable of binding oncostatin M, and also binds leukemia inhibitory factor (LIF). The receptor thus is useful for treating disorders mediated by either oncostatin
20 M or LIF.

 The gp130 may be covalently linked to the LIF-R by any suitable means, such as via a cross-linking reagent or a polypeptide linker. The gp130 and LIF-R proteins are covalently linked in a manner that does not interfere with the resulting receptor's ability to bind oncostatin M and LIF. In one embodiment of the invention, the receptor
25 is a fusion protein produced by recombinant DNA technology.

 Non-covalent bonding of gp130 to LIF-R may be achieved by any suitable means that does not interfere with the receptor's ability to bind oncostatin M and LIF. In one approach, a first compound is attached to LIF-R and a second compound that will non-covalently bond to the first compound is attached to gp130. Examples of
30 such compounds are biotin and avidin. The receptor is thus formed through the non-covalent interactions of biotin with avidin. In one embodiment of the invention, LIF-R and gp130 are recombinant polypeptides, each purified from recombinant cells and then non-covalently bonded together to form the receptor. A host cell may be transformed with two different expression vectors such that both LIF-R and gp130 are produced by
35 the recombinant host cell. LIF-R and gp130 (one or both of which are soluble fragments as described below) produced by such transformed host cells may associate to form a complex through non-covalent interactions.

PCT/US92/10272

WO 93/10151

4

"Leukemia inhibitory factor receptor" (LIF-R) refers to a protein (a cytokine receptor) that is present on the surface of various hematopoietic cells, including monocyte-macrophages and megakaryocytes, and on non-hematopoietic cells, including osteoblasts, placental trophoblasts, and liver parenchymal cells. LIF-R is capable of binding leukemia inhibitory factor (LIF) molecules and plays a role in transducing the signal provided by LIF to a cell. In the absence of any species designation, LIF-R refers generically to mammalian LIF-R, which includes, but is not limited to, human, murine, and bovine LIF-R.

The cloning of human and murine leukemia inhibitory factor receptors (LIF-R), each a single polypeptide chain, has been reported by Gearing et al. in *EMBO J.* 10:2839 (1991), which is hereby incorporated by reference in its entirety. The DNA sequence of a human LIF-R cDNA clone and the amino acid sequence encoded thereby are shown in SEQ ID NO: 5 and SEQ ID NO: 6. This cloned human cDNA encodes an N-terminal fragment of human LIF-R that includes (in order from N-terminus to C-terminus) a 44-amino acid signal sequence (amino acids -44 to -1), the entire extracellular region, a transmembrane region (the first amino acid of which is amino acid number 790 of SEQ ID NO: 5) and a portion of the cytoplasmic domain. The C-terminus of the fragment includes amino acids encoded by a poly-A segment and by a linker employed in vector construction, as described in Gearing et al., *supra*. The term "transmembrane region" as used herein refers to a string of hydrophobic amino acids positioned between the extracellular domain and the cytoplasmic domain of the protein. A plasmid vector containing the above-described cloned human LIF-R cDNA is designated pHLIFR-65 and has been deposited in *E. coli* host cells with the American Type Culture Collection on December 11, 1990 (ATCC accession no. 68491). The DNA and amino acid sequences of a full length native human LIF-R (determined by comparing the sequences of cDNA and genomic clones) have been reported by Gearing et al. *supra* and are presented herein in Figure 5.

The LIF-R encoded by the cloned cDNA (SEQ ID NO: 6) contains the entire extracellular region of LIF-R (the domain believed to be responsible for the LIF-binding activity), and binds LIF, but with lower affinity than does a naturally occurring LIF receptor found on certain normal cells. Additionally, oncostatin M competes with LIF for binding to naturally occurring high affinity LIF receptors on certain cell types (Gearing et al., *New Biologist*, 4:61, 1992) but did not bind to the above-described cloned LIF-R expressed in COS cells.

In order to investigate the possible existence of a high affinity converting subunit for the cloned single polypeptide chain LIF-R, host cells were co-transfected with the LIF-R encoding plasmid pHLIFR-65 and with pools from a human placental

WO 93/10151

PCT/US92/10272

5

cDNA library (also contained in an expression vector). The co-transfected cells were assayed for the ability to bind radiolabeled oncostatin M.

5 A positive cDNA pool was subdivided and the procedure repeated to isolate a single cDNA clone designated B10G that conferred the ability to bind oncostatin M on cells co-transfected with B10G and the LIF-R encoding plasmid pHLIFR-65. The co-transfected cells also were found to bind LIF with higher affinity than cells transfected with pHLIFR-65 alone. Host cells transfected with B10G alone exhibited low affinity oncostatin M binding sites. The B10G cloned cDNA was sequenced and found to encode a protein that is known as gp130.

10 Thus, it has now been found that a receptor comprising both LIF-R and gp130 binds LIF with higher affinity than does the single-chain LIF-R polypeptide alone. The improved LIF binding of LIF-R in combination with gp130 is described in example 1 below and depicted in Figure 1.

15 Although LIF does not bind to either high- or low-affinity oncostatin M receptors, it has now been found that oncostatin M binds to the receptors of the present invention comprising LIF-R and gp130. Oncostatin M binding is described in example 2 below and depicted in Figure 2.

20 A protein known as gp130 has been purified from cellular sources that include placental tissue and a myeloma cell line U266. A number of additional cell types have been found to express gp130 mRNA, as reported by Hibi et al., in *Cell* 63:1149 (1990). gp130 has been reported to be involved in the formation of high affinity interleukin-6 binding sites and in IL-6 signal transduction (Hibi et al. *supra*). The cloning and expression of cDNA encoding a full length gp130 protein has been reported by Hibi et al., *supra*, which is hereby incorporated by reference in its entirety.

25 The DNA and deduced amino acid sequences reported by Hibi et al. for the gp130 cloned cDNA are presented herein in figure 6. The gp130 amino acid sequence may vary from that reported by Hibi et al., e.g., leucine may be substituted for valine at position 8 in the signal sequence (numbering is as shown in Figure 6). This amino acid substitution may be attributable to genetic polymorphism (allelic variation among

30 individuals producing the protein), and results from the presence of C rather than G at nucleotide position 22.

As used herein, the term LIF-R includes variants and truncated forms of native LIF-R proteins that possess the desired LIF-binding or signal transducing activity. Likewise, the term gp130 as used herein includes variants and truncated forms of the

35 native gp130 protein that retain the desired biological activity. For gp130, the desired biological activity includes binding of oncostatin M; conferring on the inventive receptor the ability to bind oncostatin M; and increasing the affinity of the inventive

WO 93/10151

PCT/US92/10272

6

receptor for LIF. compared to the LIF binding affinity of the single-chain LIF-R polypeptide alone. Specifically included are truncated, soluble or fusion forms of LIF-R and gp130, as described below. Variants produced by adding, substituting, or deleting amino acid(s) in the native sequence are discussed in more detail below.

- 5 One example of an LIF-R polypeptide that may be employed is that encoded by the cDNA clone designated pHLIF-R-65 (SEQ ID NO: 5), as described by Gearing et al., *supra* and in example 3 below. Alternatively, a fragment comprising amino acids 1 to 945 of SEQ ID NO:5 may be employed. Amino acid 945 is the last LIF-R-specific amino acid of the polypeptide encoded by clone pHLIF-R-65, before the poly-A nucleotide segment believed to result from oligo(dT) priming at an internal site in the mRNA during preparation of the hLIF-R cDNA. (See Gearing et al., *EMBO J.*, *supra* at page 2840, column one.)

- 15 Other examples of LIF-R polypeptides that may be employed in the inventive receptors include those lacking all or part of the transmembrane region or the cytoplasmic domain of the protein. Suitable LIF-R polypeptides thus include those containing amino acids 1-x or, when the signal sequence is not desired, amino acids 45-x of the full length LIF-R sequence depicted in Figure 5, wherein x represents an integer from 833 to 1096. Amino acid number 833 is the last amino acid of the extracellular domain (i.e., before the start of the transmembrane region.) Polypeptides
- 20 terminating in amino acid number 1096 lack the last C-terminal amino acid of the full length protein. The desirability of including the signal sequence depends on such factors as the position of LIF-R in a fusion protein, as discussed below, and the intended host cells when the receptor is to be produced *via* recombinant DNA technology. Note that the numbering of amino acids in Figure 5 (taken from Gearing et al., *supra*) differs from that of SEQ ID NO: 5 because the first amino acid of the signal sequence is designated amino acid number 1 in Figure 5 but is designated -44 in SEQ ID NO: 5. Other polypeptides may be chosen with regard to sequences that are conserved in the hematopoietin receptor family. (i.e., chosen to include the boxed sequence(s) shown in Figure 5.)

- 30 One example of a suitable gp130 polypeptide is that encoded by cDNA cloned into plasmid vector pDC303 to produce a plasmid designated B10G. The source of mRNA used in producing the cDNA was human placental tissue. Plasmid B10G in *E. coli* strain DH5 α host cells was deposited with the American Type Culture Collection, Rockville, Maryland, on November 14, 1991, and assigned ATCC accession number
- 35 68827.

The DNA sequence of the gp130 cDNA contained in plasmid B10G and the amino acid sequence of the gp130 protein encoded by the cloned cDNA are presented in

WO 93/10151

PCT/US92/10272

7

SEQ ID NO: 1 and SEQ ID NO: 2. The protein comprises (in order from the N-terminus to the C-terminus) a 22-amino acid signal sequence, complete extracellular domain (amino acids 1-597), a transmembrane region (beginning with amino acid 598), and a partial cytoplasmic domain (amino acids 621-686). This truncated gp130 polypeptide differs from the equivalent portion of the Hibi et al. protein in that the eighth amino acid of the signal sequence is leucine rather than valine, as discussed above.

Another example of a suitable gp130 polypeptide comprises amino acids 1 to 496 of the SEQ ID NO: 1, which includes all of the cysteine residues found in the extracellular domain of the protein, and also contains a complete fibronectin domain. Additional examples of gp130 polypeptides are those comprising amino acids 1-298 or 98-298 of SEQ ID NO: 1.

Other gp130 polypeptides lacking all or part of the transmembrane region and/or cytoplasmic domain may be employed. Suitable gp130 polypeptides thus include those containing amino acids 1-x or, when the signal sequence is not desired, amino acids 23-x of the Figure 6 sequence, wherein x represents an integer from 619 to 917. The first amino acid of the transmembrane region is the alanine residue at position 620 in Figure 6. Polypeptides terminating at amino acid 917 lack the last C-terminal amino acid of the full length protein presented in Figure 6. Note that the numbering of amino acids in Figure 6 (taken from Hibi et al., *supra*) differs from that shown in SEQ ID NO: 1 and NO:2 because the first amino acid of the signal sequence is designated amino acid number 1 in Figure 6 but is designated -22 in SEQ ID NO: 1. Regions of the gp130 protein corresponding to domains that are conserved among certain receptors are discussed by Hibi et al., *supra*, at page 1150, column 2, and page 1151, column 1. Other truncated gp130 polypeptides chosen to include these conserved regions may be employed.

Preferred LIF-R and gp130 polypeptides are those which are soluble. In one embodiment of the present invention, the receptor comprises soluble LIF-R covalently attached to soluble gp130. "Soluble LIF-R" as used in the context of the present invention refers to polypeptides that are substantially similar in amino acid sequence to all or part of the extracellular region of a native LIF-R and that, due to the lack of a transmembrane region that would cause retention of the polypeptide on a cell membrane, are secreted upon expression. The soluble LIF-R polypeptides that may be employed retain the ability to bind LIF or, by competitively binding LIF, inhibit LIF signal transduction activity via cell surface bound LIF-R proteins. Soluble LIF-R may also include part of the transmembrane region or part of the cytoplasmic domain or other sequences, provided that the soluble LIF-R protein is capable of being secreted.

PCT/US92/10272

WO 93/10151

8

Likewise, the term "soluble gp130" as used herein refers to proteins that are substantially similar in amino acid sequence to all or part of the extracellular region of a native gp130 and are secreted upon expression but retain the desired biological activity. Soluble gp130 may include part of the transmembrane region, cytoplasmic domain, or other sequences, as long as the polypeptide is secreted.

5 Soluble LIF-R and soluble gp130 may be identified (and distinguished from their non-soluble membrane-bound counterparts) by separating intact cells which express the desired protein from the culture medium, e.g., by centrifugation, and assaying the medium (supernatant) for the presence of the desired protein. The culture
10 medium may be assayed using procedures which are similar or identical to those described in the examples below. The presence of LIF-R or gp130 in the medium indicates that the protein was secreted from the cells and thus is a soluble form of the desired protein. Soluble LIF-R and soluble gp130 may be naturally-occurring forms of these proteins. Cloning of a naturally-occurring soluble murine LIF-R is reported in
15 Gearing et al., *supra*. Alternatively, soluble fragments of LIF-R and gp130 proteins may be produced by recombinant DNA technology or otherwise isolated, as described below.

The use of soluble forms of LIF-R and gp130 is advantageous for certain applications. Purification of the proteins from recombinant host cells is facilitated,
20 since the soluble proteins are secreted from the cells. Further, soluble proteins are generally more suitable for intravenous administration and may exert their therapeutic effect (binding LIF and oncostatin M) in the bloodstream.

Soluble LIF-R polypeptides include those comprising the signal sequence and entire extracellular domain (amino acids -44 to 789 of SEQ ID NO: 5) or lacking the
25 signal sequence but containing the entire extracellular domain (amino acids 1 to 789 of SEQ ID NO: 5). Soluble gp130 polypeptides include those comprising the signal sequence and entire extracellular domain (amino acids -22 to 597 of SEQ ID NO: 1) or lacking the signal sequence but containing the entire extracellular domain (amino acids 1 to 597 of SEQ ID NO: 1). The preparation and use of these soluble polypeptides in
30 receptors of the present invention is described in examples 3-5.

Other soluble LIF-Rs are truncated upstream of the transmembrane region, but preferably include that portion of the protein that contains the residues conserved among the members of the hematopoietin receptor family (shown boxed in Figure 5), i.e., amino acids 11-479 of SEQ ID NO: 6. The N-terminus of such soluble LIF-Rs is
35 any of amino acids 1-11 (or -44 when the native signal sequence is included), and the protein extends to a C-terminus selected from any of amino acids 479 through 789. Two such soluble proteins comprise amino acids -44 - 702, 1 - 702, -44 - 775, or 1 -

WO 93/10151

PCT/US92/10272

9

755 of SEQ ID NO:6. Constructs encoding these proteins may be prepared by techniques that involve cleaving the human LIF-R cDNA of the above-described clone pHLIFR-65 (ATCC 68491) with the restriction endonucleases *Asp718* and *XmnI* or with *Asp718* and *Bsp1286I*. *Asp 718* cleaves the vector upstream of the inserted LIF-R-encoding cDNA. *XmnI* cleaves within the codon for Asp at position 702 (generating blunt ends) and *Bsp1286I* cleaves just 3' of the codon for Val at position 775 of SEQ ID NO:5. If desired, an oligonucleotide may be ligated to the 3' end of the *Asp718/Bsp1286I* fragment to extend the LIF-R sequence, e.g., through amino acid number 789. An oligonucleotide also may be ligated to the 3' end of a LIF-R fragment to add the first two amino acids of the Fc polypeptide described in example 5, and a Bgl II site useful for attaching the rest of the Fc sequence downstream of the LIF-R sequence.

Additional soluble human LIF-Rs comprise amino acids 1-678 or 1-680 of SEQ ID NO:6. When the human and murine LIF-R amino acid sequences disclosed in Gearing et al., *EMBO J.*, *supra*, are aligned (with gaps introduced to maximize identity between the two sequences), amino acid 680 of the human sequence is aligned with the last amino acid of the murine protein, and amino acid 678 is the last amino acid of the human sequence that is identical to a corresponding amino acid in the murine sequence. Since the murine protein binds LIF, the murine LIF-R contains that portion of the protein required for LIF binding.

An additional example of a soluble gp130 polypeptide comprises amino acids 22 to 582 of SEQ ID NO:2. An expression vector encoding such a protein was constructed in example 7. Soluble LIF-R and gp130 polypeptides also include those from which fibronectin type III (FNIII) domains have been deleted. From one to all of the FNIII domains may be deleted, providing the advantage of reducing the size of the protein. Preparation of such LIF-R and gp130 proteins is described in example 8.

Truncated LIF-R and gp130, including soluble polypeptides, may be prepared by any of a number of conventional techniques. In the case of recombinant proteins, a DNA fragment encoding a desired fragment may be subcloned into an expression vector. Alternatively, a desired DNA sequence may be chemically synthesized using known techniques. DNA fragments also may be produced by restriction endonuclease digestion of a full length cloned DNA sequence, and isolated by electrophoresis on agarose gels. Linkers containing restriction endonuclease cleavage site(s) may be employed to insert the desired DNA fragment into an expression vector, or the fragment may be digested at cleavage sites naturally present therein. Alternatively, proteins may be fragmented using proteolytic enzymes, for example, and the desired truncated polypeptide isolated from the digestion mixture using reversed phase HPLC.

WO 93/10151

PCT/US92/10272

10

The well known polymerase chain reaction procedure also may be employed to isolate a DNA sequence encoding a desired protein fragment. This technique is illustrated in examples 3-5 below.

5 In another approach, enzymatic treatment (e.g., using Bal 31 exonuclease) may be employed to delete terminal nucleotides from a DNA fragment to obtain a fragment having a particular desired terminus. Among the commercially available linkers are those that can be ligated to the blunt ends produced by Bal 31 digestion, and which contain restriction endonuclease cleavage site(s). Alternatively, oligonucleotides that reconstruct the N- or C- terminus of a DNA fragment to a desired point may be synthesized. The oligonucleotide may contain a restriction endonuclease cleavage site upstream of the desired coding sequence and position an initiation codon (ATG) at the N-terminus of the coding sequence.

10 The gp130 polypeptide is attached to the LIF-R polypeptide through a covalent or non-covalent linkage. Covalent attachment is preferred for certain applications, e.g. *in vivo* use, in view of the enhanced stability generally conferred by covalent, as opposed to non-covalent, bonds. In constructing the receptor of the present invention, covalent linkage may be accomplished via cross-linking reagents, polypeptide linkers, or any other suitable technique.

15 Numerous reagents useful for cross-linking one protein molecule to another are known. Heterobifunctional and homobifunctional linkers are available for this purpose from Pierce Chemical Company, Rockford, Illinois, for example. Such linkers contain two functional groups (e.g., esters and/or maleimides) that will react with certain functional groups on amino acid side chains, thus linking one polypeptide to another. The reagent and reaction conditions should be chosen such that the cross-linking does not interfere with binding of oncostatin M and LIF to the receptor.

20 One type of polypeptide linker that may be employed in the present invention separates gp130 and LIF-R domains by a distance sufficient to ensure that each domain properly folds into the secondary and tertiary structures necessary for the desired biological activity. The linker also should allow the extracellular domains of gp130 and LIF-R to assume the proper spatial orientation to form the binding site for oncostatin M and LIF. Suitable polypeptide linkers preferably (1) will adopt a flexible extended conformation, (2) will not exhibit a propensity for developing an ordered secondary structure which could interact with the functional gp130 and LIF-R domains, and (3) will have minimal hydrophobic or charged character which could promote interaction with the functional protein domains. Typical surface amino acids in flexible protein regions include Gly, Asn and Ser. Virtually any permutation of amino acid sequences containing Gly, Asn and Ser would be expected to satisfy the above criteria for a

WO 93/10151

PCT/US92/10272

11

peptide linker sequence. Other near neutral amino acids, such as Thr and Ala, may also be used in the linker sequence. Examples of such polypeptide linkers are presented below.

Another type of polypeptide linker that may be employed comprises the Fc region of an antibody. An Fc polypeptide is attached to the C-terminus of LIF-R or of the LIF-R fragment. A separate Fc polypeptide is attached to the C-terminus of gp130 or of the gp130 fragment. The two resulting polypeptide chains may be combined in a buffered solution, whereupon disulfide bonds form between the two Fc polypeptides (e.g., in the so-called hinge region, where interchain disulfide bonds are normally present in antibody molecules). Preferably, a host cell is transformed with DNA encoding both polypeptides such that the two polypeptides are co-expressed and interchain disulfide bonds form in the host cell. LIF-R is thus covalently linked to gp130 via the disulfide bonds in the linker portion of the receptor. Procedures for isolating the Fc region of an antibody are well-known and include proteolytic digestion with papain. Alternatively, an Fc polypeptide may be produced by recombinant cells or chemically synthesized. Also useful are N-terminal fragments of an antibody Fc region that contain the cysteine residues involved in disulfide bond formation at the hinge region. One example of a receptor containing an Fc polypeptide linker is illustrated in example 5 below. The receptor is depicted in Figure 4. The number and position of disulfide bonds may vary from those shown in Figure 4.

Additional examples of LIF-R/Fc and gp130/Fc fusion proteins useful in preparing receptors of the present invention are described in examples 7 and 8. Advantageously, host cells are co-transfected with two different expression vectors, one encoding soluble LIF-R/Fc and the other encoding soluble gp130/Fc. The heterodimer is believed to form intracellularly or during secretion.

Homodimers comprising two LIF-R/Fc polypeptides or two gp130/Fc polypeptides linked via disulfide bonds are also produced by certain of the transfected host cells disclosed herein. The LIF-R/Fc homodimers are useful for binding LIF and the gp130/Fc homodimers find use in binding oncostatin M. The homodimers may be separated from each other and from the heterodimer by virtue of differences in size (e.g., by gel electrophoresis). The heterodimer also may be purified by sequential immunoaffinity chromatography (described below).

In an alternative embodiment, a first fusion polypeptide comprising gp130 (or fragment thereof) upstream of an antibody light chain (or a fragment thereof) is prepared. A second fusion polypeptide comprises LIF-R upstream of an antibody heavy chain (or a heavy chain fragment, the N-terminus of which extends at least through the CH1 region. Disulfide bond(s) form between the gp130-light chain

WO 93/10151

PCT/US92/10272


12

fusion polypeptide and the LIF-R-heavy chain fusion polypeptide, thus producing a receptor of the present invention comprising a polypeptide linker. If desired, a third fusion (an LIF-R-antibody light chain fusion polypeptide) is prepared and combined with (disulfide bonded to) a fourth fusion comprising gp130 fused to an antibody heavy chain. When the two disulfide bonded molecules are combined, additional disulfide bonds form between the two Fc regions. The resulting receptor of the present invention comprising the four fusion polypeptides resembles an antibody in structure and displays the oncostatin M/LIF binding site bivalently.

A polypeptide linker may be attached to gp130 and to LIF-R by any of the conventional procedures used to attach one polypeptide to another. The cross-linking reagents available from Pierce Chemical Company as described above are among those that may be employed. Amino acids having side chains reactive with such reagents may be included in the polypeptide linker, e.g., at the termini thereof.

The gp130 and LIF-R polypeptides may be separately purified from cellular sources, and then linked together. Alternatively, the receptor of the present invention may be produced using recombinant DNA technology. The gp130 and LIF-R polypeptides may be produced separately and purified from transformed host cells for subsequent covalent linkage. In one embodiment of the present invention, a host cell is transformed/transfected with foreign DNA that encodes gp130 and LIF-R as separate polypeptides. The two polypeptides may be encoded by the same expression vector with start and stop codons for each of the two genes, or the recombinant cells may be co-transfected with two separate expression vectors. In another embodiment, the receptor is produced as a fusion protein in recombinant cells.

In one embodiment of the present invention, the receptor protein is a recombinant fusion protein of the formula:



wherein R_1 represents gp130 or a gp130 fragment; R_2 represents LIF-R or an LIF-R fragment; and L represents a polypeptide linker.

The fusion proteins of the present invention include constructs in which the C-terminal portion of gp130 is fused to the linker which is fused to the N-terminal portion of LIF-R, and also constructs in which the C-terminal portion of LIF-R is fused to the linker which is fused to the N-terminal portion of gp130. gp130 is covalently linked to LIF-R in such a manner as to produce a single protein which retains the desired biological activities of gp130 and LIF-R. The components of the fusion protein are listed in their order of occurrence (i.e., the N-terminal polypeptide is listed first, followed by the linker and then the C-terminal polypeptide).

WO 93/10151

PCT/US92/10272

13

A DNA sequence encoding a fusion protein is constructed using recombinant DNA techniques to insert separate DNA fragments encoding gp130 and LIF-R into an appropriate expression vector. The 3' end of a DNA fragment encoding gp130 is ligated (*via* the linker) to the 5' end of the DNA fragment encoding LIF-R with the reading frames of the sequences in phase to permit translation of the mRNA into a single biologically active fusion protein. Alternatively, the 3' end of a DNA fragment encoding LIF-R may be ligated (*via* the linker) to the 5' end of the DNA fragment encoding gp130, with the reading frames of the sequences in phase to permit translation of the mRNA into a single biologically active fusion protein. A DNA sequence encoding an N-terminal signal sequence may be retained on the DNA sequence encoding the N-terminal polypeptide, while stop codons, which would prevent read-through to the second (C-terminal) DNA sequence, are eliminated. Conversely, a stop codon required to end translation is retained on the second DNA sequence. DNA encoding a signal sequence is preferably removed from the DNA sequence encoding the C-terminal polypeptide.

Suitable polypeptide linkers comprise a chain of amino acids, preferably from 20 to 100 amino acids in length and most preferably from 30 to 60 amino acids in length. As discussed above, the linker advantageously comprises amino acids selected from the group consisting of glycine, asparagine, serine, threonine, and alanine. Examples of suitable polypeptide linkers include, but are not limited to, (Gly₄Ser)_n, wherein n is 4-12, preferably 8, and (Gly₄SerGly₃Ser)₂.

A DNA sequence encoding a desired polypeptide linker may be inserted between, and in the same reading frame as, the DNA sequences encoding gp130 and LIF-R using any suitable conventional technique. For example, a chemically synthesized oligonucleotide encoding the linker and containing appropriate restriction endonuclease cleavage sites may be ligated between the sequences encoding gp130 and LIF-R.

Alternatively, a chemically synthesized DNA sequence may contain a sequence complementary to the 3' terminus (without the stop codon) of either gp130 or LIF-R, followed by a linker-encoding sequence which is followed by a sequence complementary to the 5' terminus of the other of gp130 and LIF-R. Oligonucleotide directed mutagenesis is then employed to insert the linker-encoding sequence into a vector containing a direct fusion of gp130 and LIF-R.

The present invention provides an isolated DNA sequence encoding the above-described fusion protein comprising gp130, LIF-R, and a polypeptide linker, and also provides recombinant expression vectors containing the isolated DNA sequence. "Expression vector" refers to a replicable DNA construct used to express DNA which

WO 93/10151

PCT/US92/10272

14

encodes the desired protein and which includes a transcriptional unit comprising an assembly of (1) genetic element(s) having a regulatory role in gene expression, for example, promoters, operators, or enhancers, operatively linked to (2) a DNA sequence encoding a desired protein (in this case, a receptor of the present invention) which is transcribed into mRNA and translated into protein, and (3) appropriate transcription and translation initiation and termination sequences. The choice of promoter and other regulatory elements generally varies according to the intended host cell.

Proteins to be produced in yeast expression systems preferably include a leader sequence enabling extracellular secretion of translated protein by the yeast host cell. Alternatively, where recombinant protein is expressed without a leader or transport sequence, it may include an N-terminal methionine residue. This residue optionally may be subsequently cleaved from the expressed recombinant protein to provide a final product.

In the expression vectors, regulatory elements controlling transcription or translation are generally derived from mammalian, microbial, viral or insect genes. The ability to replicate in a host, usually conferred by an origin of replication, and a selection gene to facilitate recognition of transformants may additionally be incorporated. Vectors derived from retroviruses also may be employed.

DNA regions are operably linked when they are functionally related to each other. For example, DNA encoding a signal peptide (secretory leader) is operably linked to DNA for a polypeptide if the polypeptide is expressed as a precursor that is secreted through the host cell membrane; a promoter is operably linked to a coding sequence if it controls the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to permit translation. Generally, "operably linked" means contiguous and, in the case of secretory leaders, contiguous and in reading frame.

Transformed host cells are cells which have been transformed or transfected with foreign DNA using recombinant DNA techniques. In the context of the present invention, the foreign DNA includes a sequence encoding the inventive receptor. Host cells may be transformed for purposes of cloning or amplifying the foreign DNA, or may be transformed with an expression vector for production of the receptor protein. Suitable host cells for expression of the receptor include prokaryotes, yeast or higher eukaryotic cells under the control of appropriate promoters. Prokaryotes include gram negative or gram positive organisms, for example *E. coli* or bacilli. Prokaryotic expression vectors generally comprise one or more phenotypic selectable markers, for example a gene encoding proteins conferring antibiotic resistance or supplying an autotrophic requirement, and an origin of replication recognized by the host to ensure

WO 93/10151

PCT/US92/10272

15

amplification within the host. Examples of suitable prokaryotic hosts for transformation include *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium*, and various species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, although others may also be employed as a matter of choice. Higher eukaryotic cells include established cell lines of mammalian origin. Cell-free translation systems could also be employed to produce fusion protein using RNAs derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are described by Pouwels et al. (*Cloning Vectors: A Laboratory Manual*, Elsevier, New York, 1985), the relevant disclosure of which is hereby incorporated by reference.

Useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well-known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and pGEM1 (Promega Biotec, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed. *E. coli* is typically transformed using derivatives of pBR322, a plasmid derived from an *E. coli* species (Bolivar et al., *Gene* 2:95, 1977). pBR322 contains genes for ampicillin and tetracycline resistance and this provides simple means for identifying transformed cells.

Promoters commonly used in recombinant microbial expression vectors include the β -lactamase (penicillinase) and lactose promoter system (Chang et al., *Nature* 275:615, 1978; and Goeddel et al., *Nature* 281:544, 1979), the tryptophan (*trp*) promoter system (Goeddel et al., *Nucl. Acids Res.* 8:4057, 1980; and EPA 36,776) and *tac* promoter (Maniatis, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, p. 412, 1982). A particularly useful bacterial expression system employs the phage λ P_L promoter and cI857ts thermoinducible repressor. Plasmid vectors available from the American Type Culture Collection which incorporate derivatives of the λ P_L promoter include plasmid pHUB2, resident in *E. coli* strain JMB9 (ATCC 37092) and pPLc28, resident in *E. coli* RR1 (ATCC 53082).

The recombinant receptor protein may also be expressed in yeast hosts, preferably from *Saccharomyces* species, such as *S. cerevisiae*. Yeast of other genera such as *Pichia* or *Kluyveromyces* may also be employed. Yeast vectors will generally contain an origin of replication from the 2 μ m yeast plasmid or an autonomously replicating sequence (ARS), a promoter, DNA encoding the receptor fusion protein, sequences for polyadenylation and transcription termination and a selection gene. Preferably, yeast vectors will include an origin of replication and selectable markers

PCT/US92/10272

WO 93/10151

16

5 permitting transformation of both yeast and *E. coli*, e.g., the ampicillin resistance gene of *E. coli* and the *S. cerevisiae* *trp1* gene, which provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, and a promoter derived from a highly expressed yeast gene to induce transcription of a structural sequence downstream. The presence of the *trp1* lesion in the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan.

10 Suitable promoter sequences in yeast vectors include the promoters for metallothionein, 3-phosphoglycerate kinase (Hitzeman et al., *J. Biol. Chem.* 255:2073, 1980) or other glycolytic enzymes (Hess et al., *J. Adv. Enzyme Reg.* 7:149, 1968; and Holland et al., *Biochem.* 17:4900, 1978), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase and glucokinase. Suitable vectors and
15 promoters for use in yeast expression are further described in R. Hitzeman et al., EPA 73,657.

Preferred yeast vectors can be assembled using DNA sequences from pBR322 for selection and replication in *E. coli* (*Amp^r* gene and origin of replication) and yeast DNA sequences including a glucose-repressible *ADH2* promoter and α -factor secretion
20 leader. The *ADH2* promoter has been described by Russell et al. (*J. Biol. Chem.* 258:2674, 1982) and Brier et al., (*Nature* 300:724, 1982). The yeast α -factor leader, which directs secretion of heterologous proteins, can be inserted between the promoter and the structural gene to be expressed. See, e.g., Kurjan et al., *Cell* 50:922, 1982; and Brier et al., *Proc. Natl. Acad. Sci. USA* 81:5330, 1984. The leader sequence may
25 be modified to contain, near its 3' end, one or more useful restriction sites to facilitate fusion of the leader sequence to foreign genes.

Suitable yeast transformation protocols are known to those of skill in the art. An exemplary technique is described by Hinnen et al., *Proc. Natl. Acad. Sci. USA* 75:1929, (1978), selecting for *Trp^r* transformants in a selective medium consisting of
30 0.67% yeast nitrogen base, 0.5% casamino acids, 2% glucose, 10 μ g/ml adenine and 20 μ g/ml uracil.

Host strains transformed by vectors comprising the *ADH2* promoter may be grown for expression in a rich medium consisting of 1% yeast extract, 2% peptone, and 1% glucose supplemented with 80 μ g/ml adenine and 80 μ g/ml uracil.
35 Derepression of the *ADH2* promoter occurs upon exhaustion of medium glucose. Crude yeast supernatants are harvested by filtration and held at 4°C prior to further purification.

WO 93/10151

PCT/US92/10272

17

Various mammalian or insect cell culture systems can be employed to express recombinant protein. Baculovirus systems for production of heterologous proteins in insect cells are reviewed by Luckow and Summers, *Bio/Technology* 6:47 (1988). Examples of suitable mammalian host cell lines include the COS-7 lines of monkey kidney cells, described by Gluzman (*Cell* 23:175, 1981), and other cell lines capable of expressing an appropriate vector including, for examples, L cells, C127, 3T3, Chinese hamster ovary (CHO), HeLa and BHK cell lines. Mammalian expression vectors may comprise non-transcribed elements such as an origin of replication, a suitable promoter and enhancer linked to the gene to be expressed, and other 5' or 3' flanking nontranscribed sequences, and 5' or 3' nontranslated sequences, such as necessary ribosome binding sites, a poly-adenylation site, splice donor and acceptor sites, and transcriptional termination sequences.

The transcriptional and translational control sequences in expression vectors to be used in transforming vertebrate cells may be provided by viral sources. For example, commonly used promoters and enhancers are derived from Polyoma, Adenovirus 2, Simian Virus 40 (SV40), and human cytomegalovirus. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early and late promoter, enhancer, splice, and polyadenylation sites may be used to provide the other genetic elements required for expression of a heterologous DNA sequence. The early and late promoters are particularly useful because both are obtained easily from the virus as a fragment which also contains the SV40 viral origin of replication (Fiers et al., *Nature* 273:113, 1978). Smaller or larger SV40 fragments may also be used, provided the approximately 250 bp sequence extending from the Hind III site toward the BglII site located in the viral origin of replication is included. Exemplary vectors can be constructed as disclosed by Okayama and Berg (*Mol. Cell. Biol.* 3:280, 1983).

A useful system for stable high level expression of mammalian receptor cDNAs in C127 murine mammary epithelial cells can be constructed substantially as described by Cosman et al. (*Mol. Immunol.* 23:935, 1986).

Particularly preferred vectors for expression of the inventive receptor as a fusion protein are described in the examples below. The foregoing discussion is, of course, applicable to the production of recombinant fusion proteins comprising a fragment of gp130 and/or a fragment of LIF-R. Suitable fragments are discussed above, and DNA sequences encoding such fragments may be inserted into the above-described expression vectors.

The present invention provides a process for preparing the recombinant receptor of the present invention, comprising culturing a host cell transformed with an expression vector comprising a DNA sequence that encodes said receptor under

PCT/US92/10272

WO 93/10151

18

conditions that promote expression. The receptor is then purified from culture media or cell extracts.

For example, supernatants from systems which secrete recombinant protein into the culture medium can be first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate can be applied to a suitable purification matrix. For example, a suitable affinity matrix can comprise LIF or OSM. An LIF affinity matrix may be prepared by coupling recombinant human LIF to cyanogen bromide-activated Sepharose (Pharmacia) or Hydrazide Affigel (Biorad), according to manufacturer's recommendations. Sequential immunopurification using antibodies bound to a suitable support is preferred. Proteins binding to an antibody specific for LIF-R are recovered and contacted with antibody specific for gp130 on an insoluble support. Proteins immunoreactive with both antibodies may thus be identified and isolated. Alternatively, an anion exchange resin can be employed, for example, a matrix or substrate having pendant diethylaminoethyl (DEAE) groups. The matrices can be acrylamide, agarose, dextran, cellulose or other types commonly employed in protein purification. Alternatively, a cation exchange step can be employed. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or carboxymethyl groups. Sulfopropyl groups are preferred. One or more reversed-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify a fusion protein composition.

Some or all of the foregoing purification steps, in various combinations, can be employed to provide an essentially homogeneous recombinant protein. Recombinant cell culture enables the production of the fusion protein free of those contaminating proteins which may be normally associated with gp130 or LIF-R as they are found in nature in their respective species of origin, e.g., in cells, cell exudates or body fluids.

The foregoing purification procedures are among those that may be employed to purify non-recombinant receptors of the present invention as well. When linking procedures that may produce homodimers (gp130-linker-gp130 and LIF-R-linker-LIF-R) are employed, purification procedures that separate the desired heterodimer from such homodimers are employed. An example of such a procedure is sequential immunopurification as discussed above.

Recombinant protein produced in bacterial culture is usually isolated by initial extraction from cell pellets, followed by one or more concentration, salting-out, aqueous ion exchange or size exclusion chromatography steps. Finally, high performance liquid chromatography (HPLC) can be employed for final purification

WO 93/10151

PCT/US92/10272

19

steps. Microbial cells employed in expression of recombinant fusion proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

5 Fermentation of yeast which express fusion proteins as a secreted protein greatly simplifies purification. Secreted recombinant protein resulting from a large-scale fermentation can be purified by methods analogous to those disclosed by Urdal et al. (*J. Chromatog.* 296:171, 1984), involving two sequential, reversed-phase HPLC steps for purification of a recombinant protein on a preparative HPLC column.

10 The present invention also provides a pharmaceutical composition comprising a receptor protein of the present invention with a physiologically acceptable carrier or diluent. Such carriers and diluents will be nontoxic to recipients at the dosages and concentrations employed. Such compositions may, for example, comprise the receptor protein in a buffered solution, to which may be added antioxidants such as ascorbic acid, low molecular weight (less than about ten residues) polypeptides, proteins, amino acids, carbohydrates including glucose, sucrose or dextrans, chelating agents such as EDTA, glutathione and other stabilizers and excipients. The receptor of the present invention may be administered by any suitable method in a manner appropriate to the indication, such as intravenous injection, continuous infusion, sustained release from implants, etc.

20 The DNA and/or amino acid sequences of gp130 and LIF-R may vary from those presented in SEQ ID NO: 1 and SEQ ID NO: 5. Due to the known degeneracy of the genetic code, there can be considerable variation in nucleotide sequences encoding the same amino acid sequence. The DNA sequences capable of hybridizing to the native DNA sequence of SEQ ID NO: 1 or SEQ ID NO: 5 under moderately stringent conditions (50°C, 2 X SSC), and which encode a biologically active gp130 or LIF-R polypeptide, are also considered to be gp130-encoding or LIF-R-encoding DNA sequences, respectively, in the context of the present invention. Further, certain mutations in a nucleotide sequence which encodes LIF-R or gp130 will not be expressed in the final protein product. For example, nucleotide substitutions may be made to enhance expression, primarily to avoid secondary structure loops in the transcribed mRNA (see EPA 75,444A, incorporated herein by reference). Other alterations of the nucleotide sequence may be made to provide codons that are more readily translated by the selected host, e.g., the well-known *E. coli* preference codons for *E. coli* expression.

35 The amino acid sequence of native gp130 or LIF-R may be varied by substituting, deleting, adding, or inserting one or more amino acids to produce a gp130 or LIF-R variant. Variants that possess the desired biological activity of the native

WO 93/10151

PCT/US92/10272

20

gp130 and LIF-R proteins may be employed in the receptor of the present invention. Assays by which the biological activity of variant proteins may be analyzed are described in the examples below.

Alterations to the native amino acid sequence may be accomplished by any of a number of known techniques. For example, mutations can be introduced at particular loci by synthesizing oligonucleotides containing a mutant sequence, flanked by restriction sites enabling ligation to fragments of the native sequence. Following ligation, the resulting reconstructed sequence encodes an analog having the desired amino acid insertion, substitution, or deletion.

Alternatively, oligonucleotide-directed site-specific mutagenesis procedures can be employed to provide an altered gene having particular codons altered according to the substitution, deletion, or insertion required. Exemplary methods of making the alterations set forth above are disclosed by Walder et al. (*Gene* 42:133, 1986); Baner et al. (*Gene* 37:73, 1985); Craig (*BioTechniques*, January 1985, 12-19); Smith et al. (*Genetic Engineering: Principles and Methods*, Plenum Press, 1981); U.S. Patent No. 4,518,584, and U.S. Patent No. 4,737,462, which are incorporated by reference herein.

Bioequivalent variants of LIF-R and gp130 may be constructed by, for example, making various substitutions of amino acid residues or deleting terminal or internal amino acids not needed for biological activity. In one embodiment of the invention, the variant amino acid sequence is at least 80% identical, preferably at least 90% identical, to the native sequence. Percent similarity may be determined, for example, by comparing sequence information using the GAP computer program, version 6.0, available from the University of Wisconsin Genetics Computer Group (UWGCG). The GAP program utilizes the alignment method of Needleman and Wunsch (*J. Mol. Biol.* 48:443, 1970), as revised by Smith and Waterman (*Adv. Appl. Math.* 2:482, 1981). Briefly, the GAP program defines similarity as the number of aligned symbols (i.e., nucleotides or amino acids) which are similar, divided by the total number of symbols in the shorter of the two sequences. The preferred default parameters for the GAP program include: (1) a unary comparison matrix (containing a value of 1 for identities and 0 for non-identities) for nucleotides, and the weighted comparison matrix of Gribskov and Burgess, *Nucl. Acids Res.* 14:6745, 1986, as described by Schwartz and Dayhoff, eds., *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, pp. 353-358, 1979; (2) a penalty of 3.0 for each gap and an additional 0.10 penalty for each symbol in each gap; and (3) no penalty for end gaps.

WO 93/10151

PCT/US92/10272

21

Generally, substitutions should be made conservatively; i.e., the most preferred substitute amino acids are those having physiochemical characteristics resembling those of the residue to be replaced. Examples of conservative substitutions include substitution of one aliphatic residue for another, such as Ile, Val, Leu, or Ala for one another, or substitutions of one polar residue for another, such as between Lys and Arg; Glu and Asp; or Gln and Asn. Other such conservative substitutions, for example, substitutions of entire regions having similar hydrophobicity characteristics, are well known. Moreover, particular amino acid differences between human, murine and other mammalian LIF-Rs is suggestive of additional conservative substitutions that may be made without altering the essential biological characteristics of LIF-R.

Cysteine residues can be deleted or replaced with other amino acids to prevent formation of unnecessary or incorrect intramolecular disulfide bridges upon renaturation. Adjacent dibasic amino acid residues may be modified to enhance expression in yeast systems in which KEX2 protease activity is present.

EP212,914 discloses the use of site-specific mutagenesis to inactivate KEX2 protease processing sites in a protein. KEX2 protease processing sites are inactivated by deleting, adding or substituting residues to alter Arg-Arg, Arg-Lys, and Lys-Arg pairs to eliminate the occurrence of these adjacent basic residues. Lys-Lys pairings are considerably less susceptible to KEX2 cleavage, and conversion of Arg-Lys or Lys-Arg to Lys-Lys represents a conservative and preferred approach to inactivating KEX2 sites.

Hydrophilic amino acids may be substituted for hydrophobic amino acids in the transmembrane region and/or intracellular domain of gp130 and LIF-R to enhance water solubility of the proteins. Addition of amino acids to the native sequence may result from translation of in-frame codons present in linkers used in constructing cloning or expression vectors. The LIF-R encoded by clone pHLIF-R-65 contains such linker-encoded amino acids at the C-terminus, as described by Gearing et al., *supra*.

The present invention also includes proteins with or without associated native-pattern glycosylation. Expression of DNAs encoding the fusion proteins in bacteria such as *E. coli* provides non-glycosylated molecules. Functional mutant analogs having inactivated N-glycosylation sites can be produced by oligonucleotide synthesis and ligation or by site-specific mutagenesis techniques. These analog proteins can be produced in a homogeneous, reduced-carbohydrate form in good yield using yeast expression systems. N-glycosylation sites in eukaryotic proteins are characterized by the amino acid triplet Asn-A1-Z, where A1 is any amino acid except Pro, and Z is Ser or Thr. In this sequence, asparagine provides a side chain amino group for covalent

WO 93/10151

PCT/US92/10272

22

attachment of carbohydrate. Such a site can be eliminated by substituting another amino acid for Asn or for residue Z, deleting Asn or Z, or inserting a non-Z amino acid between A₁ and Z, or an amino acid other than Asn between Asn and A₁. Known procedures for inactivating N-glycosylation sites in proteins include those described in U.S. Patent 5,071,972 and EP 276,846.

5 Variants of the receptor proteins of the present invention also include various structural forms of the primary protein which retain biological activity. Due to the presence of ionizable amino and carboxyl groups, for example, a receptor protein may be in the form of acidic or basic salts, or may be in neutral form. Individual amino acid residues may also be modified by oxidation or reduction.

10 The primary amino acid structure also may be modified by forming covalent or aggregative conjugates with other chemical moieties, such as glycosyl groups, lipids, phosphate, acetyl groups and the like. Covalent derivatives are prepared by linking particular functional groups to amino acid side chains or at the N- or C-termini. Other derivatives of the receptor protein within the scope of this invention include covalent or aggregative conjugates of the receptor protein with other proteins or polypeptides, such as by synthesis in recombinant culture as N- or C-terminal fusions. For example, the conjugated polypeptide may be a signal (or leader) polypeptide sequence at the N-terminal region of the protein which co-translationally or post-translationally directs transfer of the protein from its site of synthesis to its site of function inside or outside of the cell membrane or wall (e.g., the yeast α -factor leader). Peptides may also be added to facilitate purification or identification of the fusion protein (e.g., poly-His). The amino acid sequence of the fusion protein can also be linked to the peptide Asp-Tyr-Lys-Asp-Asp-Asp-Asp-Lys (DYKDDDDK) (Hopp et al., *BioTechnology* 6:1204, 1988). The latter sequence is highly antigenic and provides an epitope reversibly bound by a specific monoclonal antibody, enabling rapid assay and facile purification of expressed recombinant protein. This sequence is also specifically cleaved by bovine mucosal enterokinase at the residue immediately following the Asp-Lys pairing. Receptor proteins capped with this peptide may also be resistant to intracellular degradation in *E. coli*.

30 The receptors of the present invention are primarily useful as oncostatin M binding reagents, and may be administered *in vivo* to inhibit a biological activity of oncostatin M (including signal transduction). The inventive receptors also have use as LIF binding reagents.

95 Disorders mediated by either oncostatin M or LIF may be treated by administering a therapeutically effective amount of the receptor of the present invention to a human or mammalian patient afflicted with such a disorder. A disorder is said to

WO 93/10151

PCT/US92/10272

23

be mediated by oncostatin M or LIF when biologically active oncostatin M or LIF causes (directly or indirectly) or exacerbates the disorder. Soluble receptor proteins can be used to competitively bind to LIF and oncostatin M, thereby inhibiting binding of LIF and oncostatin M to cell surface receptors.

5 As discussed in example 2, gp130 has now been found to bind oncostatin M, although with lower affinity than the inventive receptors comprising both gp130 and LIF-R. gp130 may be administered to treat conditions mediated by oncostatin M, although a gp130/LIF-R receptor of the present invention would be preferred for such a purpose.

10 Oncostatin M has been reported to stimulate hematopoiesis, stimulate epithelial cell proliferation, increase plasmin activity (thereby inducing fibrinolysis), inhibit angiogenesis and suppress expression of major histocompatibility complex antigens on endothelial cells. See PCT application WO 9109057 and European patent application no. 422,186. When these or other biological effects of oncostatin M are undesirable, a
15 receptor of the present invention may be administered to bind oncostatin M.

Oncostatin M is believed to stimulate production of the cytokine interleukin-6 (IL-6), as reported by Brown et al., *J. Immunol.* 147:2175 (1991). Oncostatin M therefore indirectly mediates disorders associated with the presence of IL-6. IL-6 has been reported to be involved in the pathogenesis of AIDS-associated Kaposi's sarcoma (deWit et al., *J. Intern. Med. [England]* 229:539 [1991]). Binding of oncostatin M by
20 a receptor of the present invention thus may be useful in treating Kaposi's sarcoma. Alternatively, but less preferably, gp130 may be administered to treat Kaposi's sarcoma.

Among the disorders mediated by LIF are lipoprotein metabolism defects such
25 as atherosclerosis and obesity, as well as disorders of bone and calcium metabolism or disorders associated with LIF overproduction that affect hepatocytes, neurons, or leukocytes. The regulation of embryonic and hematopoietic stem cells by LIF may also be manipulated with the receptor. A soluble form of the receptor may also be used to treat leukemic cells which respond to LIF by proliferating. LIF also may play a role in
30 inducing cachexia in cancer or AIDS patients. The receptor, or antibodies thereto, may also be useful as a diagnostic reagent to detect diseases characterized by the presence of abnormal LIF-R.

Oncostatin M and LIF are different proteins, but share certain structural and biological properties. If inhibition of a biological activity shared by oncostatin M and
35 LIF is desired, the receptor of the present invention offers the benefit of binding both of these proteins exhibiting the particular biological activity. A receptor binding only one

WO 93/10151

PCT/US92/10272

24

of the proteins would leave the other protein active and continuing to mediate the disorder.

Receptor proteins or derivatives thereof may also be used as reagents in receptor-based immunoassays, reagents in assays for oncostatin M or LIF, or as binding agents for affinity purification of oncostatin M or LIF. The receptor proteins of the present invention may be used as immunogens in conventional procedures for production of polyclonal or monoclonal antibodies. Such antibodies may be employed on immunoaffinity columns for purification of the receptor, or as components of diagnostic or research assays. Derivatives may also be obtained by attaching additional polypeptide(s), e.g., by using a cross-linking agent, such as N-maleimidobenzoyl succinimide ester that reacts with cysteine and lysine residues. Receptor proteins may also be covalently bound through reactive side groups to various insoluble substrates, such as cyanogen bromide-activated, bisoxirane-activated, carbonyldiimidazole-activated or tosyl-activated agarose structures, or by adsorbing to polyolefin surfaces (with or without glutaraldehyde cross-linking).

The following examples are provided to illustrate certain embodiments of the invention, and are not to be construed as limiting the scope of the invention.

EXAMPLES

20

Example 1

Assay to Detect Binding of LIF

Recombinant human LIF is expressed in yeast and purified to homogeneity essentially as described by Hopp, et al., *Bio/Technology* 6:1204 (1988). The purified protein is radiolabeled using a commercially available enzymobead radioiodination reagent (BioRad). In this procedure 10 µg LIF in 50 µl 0.2 M sodium phosphate, pH 7.2, are combined with 50 µl enzymobead reagent, 2 mCi of sodium iodide in 20 µl of 0.05 M sodium phosphate pH 7.0 and 10 µl of 2.5% 8-D-glucose. After 10 minutes at 25°C, sodium azide (20 µl of 50 mM) and sodium metabisulfite (10 µl of 5 mg/ml) are added and incubation is continued for 5 min. at 25°C. The reaction mixture is fractionated by gel filtration on a 2 ml bed volume of Sephadex® G-25 (Sigma) equilibrated in Roswell Park Memorial Institute (RPMI) 1640 medium containing 2.5% (w/v) bovine serum albumin (BSA), 0.2% (w/v) sodium azide and 20 mM Hepes pH 7.4 (binding medium). The final pool of ¹²⁵I-LIF is diluted to a working stock solution of 3×10^{-8} M in binding medium and stored for up to one month at 4°C

WO-93/10151

PCT/US92/10272

25

without detectable loss of receptor binding activity. The specific activity is routinely in the range of $6-8 \times 10^{15}$ cpm/nmole LIF.

5 The radiolabeled LIF may be employed in any of a number of conventional assay procedures to determine whether a given protein or cell binds LIF. Examples of such assays are those that detect binding of the radiolabeled LIF to cells expressing an LIF-binding protein on the cell surface. The radiolabeled LIF also may be employed in assays for the presence of LIF-binding proteins in cell culture medium (e.g. LIF-binding proteins secreted by recombinant cells). Proteins in cell extracts (e.g. from recombinant cells) also may be assayed for the ability to bind the radiolabeled LIF.

10 In one assay procedure, cells transformed/transfected with an expression system encoding a protein to be tested for ability to bind LIF are plated at a density of 2×10^5 cells/well in either 6 well dishes (Falcon) or single well chambered slides (Lab-Tek). Both dishes and slides are pretreated with 1 ml human fibronectin (10 μ g/ml in PBS) for 30 minutes followed by 1 wash with PBS. After 48 to 72 hours, cells are
15 assayed for expression of LIF binding proteins by binding radioiodinated LIF using the following slide autoradiography technique. Transfected cells are washed once with binding medium (RPMI media 1640 containing 25 mg/ml bovine serum albumin (BSA), 2 mg/ml sodium azide, 20 mM HEPES, pH 7.2, and 50 mg/ml nonfat dry milk (NFDM) and incubated for 2 hours at 4°C with 1 ml binding medium + NFDM
20 containing 1.25×10^{-9} M 125 I-LIF. After incubation, cells in the chambered slides are washed three times with binding buffer + NFDM, followed by 2 washes with PBS, pH 7.3, to remove unbound 125 I-LIF. The cells are fixed by incubating for 30 minutes at room temperature in 10% glutaraldehyde in PBS, pH 7.3, washed twice in PBS, and air dried. The slides are dipped in Kodak NTB-2 photographic emulsion (5x dilution in
25 water) and exposed in the dark for 12 hours to 7 days at 4°C in a light proof box. The slides are then developed for approximately 5 minutes in Kodak D19 developer (40 g/500 ml water), rinsed in water and fixed in Agfa G433C fixer. The slides are individually examined with a microscope at 25-40x magnification and positive cells that bind LIF are identified by the presence of autoradiographic silver grains against a light
30 background.

Cells in the 6 well plates are washed once with binding buffer + NFDM followed by 3 washings with PBS, pH 7.3, to remove unbound 125 I-LIF. The cells are then trypsinized to remove them from the plate and bound 125 I-LIF is counted on a gamma counter.

35 The cells in transfectant pool(s) testing positive are subdivided into smaller pools and the screening process is repeated (with further subdividing of the pools as necessary) until an individual clone expressing LIF-binding protein is isolated. Non-

WO 93/10151

PCT/US92/10272

26

specific binding of ^{125}I -LIF may be measured in the presence of 200-fold or greater excess of unlabeled LIF. As a control, the same host cells transfected with a vector lacking LIF-R-encoding sequences should be assayed to determine whether background endogenous LIF receptors are present on the host cells.

5 In another assay procedure, cells producing a soluble LIF-binding protein that is released from the cells into the culture medium may be identified. Cells are collected by centrifugation from a culture broth. The supernatant (culture medium) is concentrated 10-fold; and 1 μl aliquots are spotted onto nitrocellulose filters and allowed to air dry. Additional binding sites are blocked by overnight incubation at 4°C
10 in the above-described binding medium containing 3% non-fat dry milk (BMNFDM). Filters are incubated for 2h at 4°C in BMNFDM containing 1 nM ^{125}I -LIF in the presence or absence of 200 nM unlabeled LIF, then washed (3 x 5 min) in PBS. Filters are exposed to photographic film for 48 hr at room temperature.

The results of one LIF binding assay conducted according to the following
15 procedure are shown in Figure 1. Host cells transfected with vector(s) encoding LIF-R or gp130 as described below were assayed for the ability to bind LIF. The host cells were the monkey kidney cell line designated COS-7, described by Gluzman, *Cell* 23:175 (1981). In separate transfections, COS-7 cells were transfected with the following combinations of vectors. The different types of transfected cells (and non-
20 transfected control cells) are designated A-F as shown below, and the curves representing the LIF-binding assay data for each transfected or control cell type are also labeled A-F in Figure 1.

(A) B10G (the gp130 encoding vector described in example 3) and
25 pHLIFR-65 (the LIF-R encoding vector described in example 3)

(B) pHLIFR-65 and control vector CAV (a control vector that does not
encode LIF-R or gp130; controls for plasmid dilution so that results can be more
accurately compared with those of COS-7 cells co-transfected with both a gp130
30 encoding vector and an LIF-R encoding vector)

(C) B10G and pHLIFR-65; transfected cells were preincubated with
non-radiolabeled oncostatin M before incubation with ^{125}I -LIF

(D) pHLIFR-65 and CAV; transfected cells were preincubated with non-
35 radiolabeled oncostatin M before incubation with ^{125}I -LIF

WO 93/10151

PCT/US92/10272

27

(E) non-transfected COS-7 cells (control)

(F) B10G and CAV

5 The assay was performed by a phthalate oil separation method essentially as described by Dower et al., *J. Immunol.* 132:751 (1984) and Park et al., *J. Biol. Chem.* 261:4177 (1986). Briefly, the COS-7 host cells were released from 10 cm tissue culture plates two days after transfection by incubation in non-enzymatic cell dissociation buffer (Sigma) at 37°C for 30-60 minutes. Cells were then washed with
10 the above-described binding medium and resuspended in binding medium at 5×10^6 cells/ml. 50 μ l aliquots of the cells were incubated with serial dilutions of 125 I-LIF at room temperature for one hour with agitation (in the presence or absence of a 200-fold excess of unlabeled LIF) in a total volume of 150 μ l. The unlabeled LIF allowed for calculation of the non-specific background binding of LIF. Duplicate aliquots (60 μ l) of
15 each incubation mixture were then transferred to a polyethylene centrifuge tube containing a phthalate oil mixture comprising 1.5 parts dibutylphthalate to 1 part bis(s-ethylhexyl)phthalate.

The cells were separated from unbound 125 I-LIF by centrifugation for five minutes at 15,000 x g in an Eppendorf microfuge. The centrifuge tubes were cut to
20 separate the pellet of cells (containing bound 125 I-LIF) from the supernatant containing unbound 125 I-LIF. The radioactivity in both parts was then determined on a gamma counter. The determinations of both cell-bound and unbound radioactivity from the two 60 μ l aliquots were averaged for subsequent calculations.

The results are presented in Figure 1 as standard Scatchard transformations of
25 the biological data. The data are reported as the ratio of molecules of 125 I-LIF bound per cell, to free 125 I-LIF molecules (y-axis) versus molecules of 125 I-LIF bound per cell (x-axis). The dissociation constants (K_D) are shown in Figure 1, along with the number of LIF-binding sites per cell. Since a saturating amount of radiolabeled LIF was offered, the number of molecules of radiolabeled LIF bound per cell is considered
30 equivalent to the number of LIF binding sites per cell.

As shown by curve A of Figure 1, COS-7 cells co-transfected with a gp130 encoding vector (B10G) and an LIF-R encoding vector (pHLIFR-65) demonstrated high affinity LIF binding ($K_D = 9 \times 10^{-10}$ M). When these same co-transfected COS-7 cells were preincubated with non-radiolabeled oncostatin M before incubation with
35 125 I-LIF (curve C), binding of LIF was greatly reduced ($K_D = 4.2 \times 10^{-9}$ M). Oncostatin M thus competes with LIF for binding sites on these transfected cells.

WO 93/10151

PCT/US92/10272

28

COS-7 cells transfected with a vector encoding the single-polypeptide chain LIF-R (pHLIFR-65) and with the control vector CAV bound LIF (curve B; $K_D=2.4 \times 10^{-9}M$), but with lower affinity than the cells producing both gp130 and LIF-R. The COS-7 cells display endogenous high affinity simian LIF receptors (curve E; K_D about $3 \times 10^{-11}M$). Transfection with pHLIFR-65 (encoding the single polypeptide LIF-R) results in display of additional low affinity LIF receptors ($K_D=2.4 \times 10^{-9}M$; curve B, site 2) as well as the simian LIF receptors $K_D=3.3 \times 10^{-11}M$; curve B, site 1).

When the COS-7 cells transfected with pHLIFR-65 and CAV were preincubated with non-radiolabeled oncostatin M before incubation with ^{125}I -LIF (CURVE D), binding of LIF to the LIF-R expressed by pHLIFR-65 was essentially unchanged compared to the same transfected cells not preincubated with oncostatin M. Oncostatin M thus does not compete with LIF for binding to the single polypeptide chain LIF-R. However, the binding of LIF to the endogenous simian high affinity LIF-R on the COS-7 cells was competed.

The COS-7 cells co-transfected with the gp130 encoding vector and the CAV control vector (curve F) did not bind LIF in any measurable amount above the amount of binding to the non-transfected COS-7 cells (curve E).

Example 2

Assay to Detect Binding of Oncostatin M

20

Oncostatin M may be purified from cells in which the protein is naturally found, or from cells transformed with an expression vector encoding oncostatin M. One source of oncostatin M is phorbol ester-treated U937 cells, as described by Zarling et al., *PNAS U.S.A.* 83:9739 (1986). Purification of recombinant oncostatin M is described by Linsley et al., *J. Biol. Chem.* 264:4282-4289 (1989), which is hereby incorporated by reference in its entirety.

Preferrably, oncostatin M is produced in yeast cells transformed with a suitable expression vector. A DNA sequence encoding a signal sequence (e.g., a yeast alpha-factor leader sequence) may be fused to the N-terminus of the oncostatin M encoding DNA sequence to promote secretion of the protein from the host cells. The protein when initially produced may also comprise an N-terminal identification leader (e.g., a "flag" sequence such as Asp-Tyr-Lys-Asp⁴-Lys) as described by Hopp et al., *Bio/Technology* 6:1204 (1988). The flag sequence is highly antigenic and provides an epitope reversibly bound by a specific monoclonal antibody, enabling facile purification of the expressed recombinant protein. This sequence is also specifically cleaved by bovine mucosal enterokinase at the residue immediately following the Asp-Lys pairing.

WO 93/10151

PCT/US92/10272

29

Neither the signal sequence nor the flag sequence is found on the processed final oncostatin M product.

Oncostatin M may be radiolabeled using any suitable conventional procedure, such as the radioiodination procedure employed to radiolabel LIF in Example 1. The radio-iodination of oncostatin M has also been described by Linsley et al., *supra*.

The resulting radiolabeled oncostatin M may be substituted for radiolabeled LIF (using the same concentrations and other reaction parameters) in the assay procedures described in Example 1 in order to detect proteins and cells that bind oncostatin M. An assay for binding of ^{125}I -oncostatin M to cells is also described in Linsley et al., *supra*.

The results of one oncostatin M binding assay are shown in Figure 2. COS-7 cells transfected with vector(s) encoding gp130 or LIF-R were assayed for the ability to bind oncostatin M. In separate transfections, COS-7 cells were transfected with the following combinations of vectors. The different types of transfected cells (and non-transfected control cells) are designated A-E as shown below, and the corresponding curves representing the oncostatin M binding assay data for each cell type are also labeled A-E in Figure 2.

(A) B10G (the gp130 encoding vector described in example 3) and pHLIFR-65 (the LIF-R encoding vector described in example 3)

(B) B10G and pHLIFR-65; transfected cells were preincubated with non-radiolabeled LIF before incubation with ^{125}I -oncostatin M

(C) pHLIFR-65 and CAV (a control vector that does not encode LIF-R or gp130; controls for plasmid dilution so that results can be more accurately compared with those of COS-7 cells co-transfected with both a gp130 encoding vector and an LIF-R encoding vector)

(D) non-transfected COS-7 cells (control)

(E) B10G and CAV

The assay was performed by the phthalate oil separation method described in example 1 (but substituting oncostatin M for LIF). The results are presented in Figure 2 as standard Scatchard transformations of the biological data. The data are reported as the ratio of molecules of ^{125}I -oncostatin M bound per cell, to free ^{125}I -oncostatin M

WO 93/10151

PCT/US92/10272

30

molecules (y-axis) versus molecules of ^{125}I -oncostatin M bound per cell (x-axis). The dissociation constants (K_D) are shown in Figure 2, along with the number of oncostatin M-binding sites per cell. Since a saturating amount of radiolabeled oncostatin M was offered, the number of molecules of radiolabeled oncostatin M bound per cell is considered equivalent to the number of oncostatin M binding sites per cell.

As shown by curve A in Figure 2, COS-7 cells co-transfected with a gp130 encoding vector (B10G) and an LIF-R encoding vector (pHLIFR-65) demonstrated the ability to bind oncostatin M with high affinity ($K_D=2.4 \times 10^{-10}\text{M}$).

COS-7 cells co-transfected with a vector encoding the single-polypeptide chain LIF-R (pHLIFR-65) and with the control vector CAV (curve C) did not bind oncostatin M in any significant amount above that bound by the non-transfected COS-7 cells (curve D).

COS-7 cells co-transfected with pHLIFR-65 and B10G and preincubated with non-radiolabeled LIF before incubation with ^{125}I -oncostatin M (curve B) did not bind oncostatin M in any measurable amount above that bound by the non-transfected COS-7 cells. LIF thus competes with oncostatin M for binding sites on the recombinant cells.

The experimental conditions of this assay (the results of which are shown in Figure 2) were not appropriate for accurate detection of low affinity oncostatin M receptors. Thus, a separate experiment (phthalate oil separation method) was conducted to compare oncostatin M binding by COS-7 cells transfected with B10G alone (no CAV control vector) with oncostatin M binding to non-transfected COS-7 cells. Non-transfected COS-7 cells assayed as a control demonstrated a small number of high affinity oncostatin M receptors ($K_D=3.6 \times 10^{-10}\text{M}$). The cells transfected with B10G demonstrated additional low affinity binding of oncostatin M ($K_D=7.7 \times 10^{-9}\text{M}$). The results of this oncostatin M binding assay are shown in Figure 3 as Scatchard transformations of the biological data. The data are reported as the ratio of molecules of ^{125}I -oncostatin M bound per cell, to free ^{125}I -oncostatin M molecules (y-axis) versus molecules of ^{125}I -oncostatin M bound per cell (x-axis). The scale in Figure 3 differs from that of Figures 1 and 2 so that the difference in oncostatin M binding by the gp130-producing cells compared to the control cells can be more readily visualized.

Disorders mediated by oncostatin M thus may be treated by administering gp130 or a fragment thereof. Receptors comprising both gp130 and LIF-R are preferred for use in treating such conditions, however, in view of the higher affinity of such receptors for oncostatin M compared to the affinity of gp130 alone for oncostatin M. gp130 also may be employed as an oncostatin M-binding reagent in diagnostic and research assays.

WO 93/10151

PCT/US92/10272

31

Example 3

Preparation of a Recombinant Fusion Protein Designated LIF-R-Linker-gp130

5 A recombinant receptor protein of the present invention is prepared by the following procedure. The receptor comprises an LIF-R fragment at the N-terminus anached to a gp130 fragment through a polypeptide linker. The polypeptide linker is of the formula (Gly₄Ser)₈. An oligonucleotide encoding a portion of the linker sequence, i.e., the sequence Ser(Gly₄Ser)₆Gly is synthesized by any of the conventional known
 10 procedures for oligonucleotide synthesis. The DNA and encoded amino acids sequences of the double-stranded oligonucleotide are as follows:

SEQ ID NO: 7

15 Bam HI

5' GATCC GGT GGA GGT GGT TCT GGT GGA GGT GGT TCA GGT GGT GGA GGA TCA
 3' G CCT CCT CCA CCA AGA CCA CCT CCA CCA AGT CCA CCA CCT CCT AGT
 Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser

20

BspMII XbaI

25 GGA GGT GGT GGA TCA GGT GGA GGA GGT TCT CCA GGT GGA CCT TCC GGA T 3'
 CCT CCA CCA CCT AGT CCA CCT CCT CCA AGA CCT CCA CCT CCA AGG CCT AGATC
 5' Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly

30 The remaining portion of the linker is added during vector construction as described below. This oligonucleotide as well as those discussed below may be synthesized on an automated DNA synthesis machine such as those available from Biosearch, Inc., San Rafael, California or Applied Biosystems.

35 The linker encoding oligonucleotide is cloned into a vector that preferably contains multiple restriction endonuclease cleavage sites that may be employed for inserting the sequences encoding LIF-R and gp130 on either side of, and in the same reading frame as, the sequence encoding the linker. One such vector is designated pBLUESCRIPT SK[®] which is available from Stratagene Cloning Systems, La Jolla, California. This plasmid vector is replicable in *E. coli* and contains a polylinker segment that includes 21 unique restriction sites. The plasmid is digested with the
 40 restriction enzymes BamHI and XbaI and the linker-encoding oligonucleotide is ligated into the vector using conventional techniques. A recombinant vector containing the inserted oligonucleotide sequence is identified by restriction endonuclease analysis and

PCT/US92/10272

WO 93/10151

32

sizing by gel electrophoresis. A DNA sequence encoding LIF-R is inserted into the pBLUESCRIPT SK[®] vector upstream of the linker- encoding oligonucleotide and a DNA sequence encoding gp130 is inserted downstream of the linker sequence. cDNA molecules encoding soluble fragments of LIF-R and gp130 were isolated and amplified using the well known polymerase chain reaction (PCR) procedure. The following oligonucleotides were synthesized for use in the PCR procedures:

SEQ ID NO: 8 (Oligonucleotide No. 1)

10 5' SalI
GATATGTCGACGATGATGGATAITTTACGTATGTTTG 3'

15 SEQ ID NO: 9 (Oligonucleotide No. 2)

3' CATACATACACCACTGTTTCCTTTTAAAGACCTCCTCCACCTAGGTACG 5'
BamHI

20 SEQ ID NO: 10 (Oligonucleotide No. 3)

25 5' BspMI
CGCGTCCGGAGGAGGTGGATCTGAACTTCTAGATCCATGTGGTTATATC 3'

30 SEQ ID No. 11 (Oligonucleotide No. 4)

3' CAAACGAGTTCCTCTTTAACTTATCCGCCGGCGTACG 5'

Oligonucleotides 1 and 2 are used in a PCR reaction to isolate a soluble fragment of LIF-R. The template employed in the reaction is the human LIF-R cDNA cloned as described by Gearing et al. *supra*. The DNA and encoded amino acid sequences of the cDNA clone are represented in SEQ ID NO: 5. The cloning vector which contains this human LIF-R cDNA clone was deposited in *E. coli* host cells with the American Type Culture Collection, Rockville, Maryland, U.S.A. on December 11, 1990, under the name pHLIFR-65 (ATCC Accession Number 68491). The deposit was made under the conditions of the Budapest Treaty. The 5' primer is oligonucleotide No. 1, which includes a DNA sequence encoding the first 8 amino acids of the signal sequence of LIF-R and also comprises upstream sequences that introduce a Sal I restriction endonuclease cleavage site. Oligonucleotide No. 1 is capable of annealing to the (-) strand that is complementary to nucleotides 179-202 of SEQ ID NO: 5. The 3' primer is oligonucleotide No. 2, which contains a sequence

WO 93/10151

PCT/US92/10272

33

complementary to nucleotides 2651-2677 of SEQ ID NO: 5 (i.e., includes anti-sense nucleotides encoding the last nine amino acids of the extracellular domain of LIF-R). Immediately downstream of the LIF-R encoding sequence, oligonucleotide No. 2 contains a sequence encoding (Gly)₄ Ser, and also introduces a BamHI restriction
5 endonuclease cleavage site.

A PCR reaction employing oligonucleotides Nos. 1 and 2 thus isolates and amplifies a DNA sequence encoding an LIF-R fragment containing the entire signal sequence and the entire extracellular domain but lacking the transmembrane region and the extracellular domain. The (Gly)₄ Ser sequence attached to the 3' terminus of the
10 LIF-R fragment is part of the polypeptide linker in the final construct.

Any suitable PCR procedure may be employed. One such procedure is described in Sarki et al., *Science* 239:487 (1988). Another is described in *Recombinant DNA Methodology*, Wu et al., eds., Academic Press Inc., San Diego (1989), pp. 189-196. In general, PCR reactions involve combining the 5' and 3'
15 nucleotides with the template DNA and each of the four deoxynucleoside triphosphates in a suitable buffered solution. The solution is heated, (e.g. from 95° to 100°C) to denature the double-stranded DNA template and is then cooled before addition of a DNA polymerase enzyme. Multiple cycles of the reactions are carried out in order to amplify the desired DNA fragment.

An example of a suitable PCR procedure is as follows. All temperatures are in degrees centigrade. The following PCR reagents are added to a 1.5 ml Eppendorf microfuge tube: 10 µl of 10X PCR buffer (500 mM KCl, 100 mM Tris-HCl, pH 8.3 at 25°C, 25 mM MgCl₂, and 1 mg/ml gelatin) (Perkin-Elmer Cetus, Norwalk, CN), 10 µl of a 2 mM solution containing each dNTP (2 mM dATP, 2 mM dCTP, 2 mM dGTP and 2 mM dTTP), 2.5 units (0.5 µl of standard 5000 units/ml solution) of *Taq* DNA
25 polymerase (Perkin-Elmer Cetus), 50 ng of template DNA, 5 µl of a 20 µM solution of each of oligonucleotide primers 1 and 2, and 74.5 µl water to a final volume of 100 µl. The final mixture is then overlaid with 100 µl paraffin oil. PCR is carried out using a DNA thermal cycler (Ericomp, San Diego, CA) by initially denaturing the template at
30 94° for 90 seconds, reannealing at 55° for 75 seconds and extending the cDNA at 72° for 150 seconds. PCR is carried out for an additional 20 cycles of amplification using a step program (denaturation at 94°, 25 sec; annealing at 55°, 45 sec; extension at 72°, 150 sec.), followed by a 5 minute extension at 72°.

The sample is removed from the paraffin oil and DNA extracted by
35 phenolchloroform extraction and spun column chromatography over G-50 (Boehringer Mannheim). A 10 µl aliquot of the extracted DNA is separated by electrophoresis on

WO 93/10151

PCI/US92/10272

34

1% SeaKem™ agarose (FMC BioProducts, Rockland, ME) and stained with ethidium bromide to confirm that the DNA fragment size is consistent with the predicted product.

5 The PCR-amplified cDNA products are then digested with SalI and BamHI restriction enzymes using standard procedures. The SalI/BamHI restriction fragment is then separated by gel electrophoresis, e.g., on a 1.2% SeaPlaque™ low gelling temperature (LGT) agarose, and the band representing the desired fragment is isolated. The fragment is inserted into a vector encoding the desired fusion protein as described below.

10 A plasmid vector containing human gp130 cDNA was deposited in *E. coli* strain DH5α host cells with the American Type Culture Collection, Rockville, Maryland under the name B10G/pDC303 (DH5α) on November 14, 1991 and assigned ATCC Accession No. 68827. The deposit was made under the conditions of the Budapest Treaty. The DNA and encoded amino acid sequences of this cloned cDNA are shown in SEQ ID NO: 1.

15 Oligonucleotides 3 and 4 are employed in the polymerase chain reaction procedure to amplify and isolate a DNA fragment encoding Ser(Gly)₄ Ser followed by amino acids 1 to 597 of SEQ ID NO: 1 (the entire extracellular domain of the mature gp130 protein). The 5' primer, oligonucleotide No. 3, includes nucleotides 310 to 336 of SEQ ID NO: 1, which encode the first nine amino acids of the mature gp130 protein. This nucleotide sequence is capable of annealing to the (-) strand that is
20 complementary to nucleotides 310 to 336 of SEQ ID NO: 1. Oligonucleotide No. 3 also encodes a Ser(Gly)₄ Ser sequence directly upstream of (and in the same reading frame as) the gp130 sequence, and further positions a BspMII restriction endonuclease cleavage site near the 5' terminus of the Ser(Gly)₄ Ser-encoding sequence.

25 The 3' primer, oligonucleotide No. 4, includes a sequence complementary to nucleotides 2080 to 2100 of SEQ ID NO: 1, i.e., includes anti-sense nucleotides encoding the last seven amino acids of the gp130 extracellular domain. Oligonucleotide No. 4 positions a stop codon immediately after the gp130 sequence and also inserts a
30 NotI restriction site downstream. Following amplification of the gp130 fragment by PCR, the PCR reaction products are digested with BspMII and NotI and the desired fragment is isolated.

The above-described LIF-R, Ser(Gly)₄Ser)₆Gly linker, and gp130 encoding fragments are assembled into a single DNA sequence as follows. The Ser(Gly)₄Ser)₆Gly linker fragment is excised from the pBLUESCRIPT SK® vector by
35 digestion with BamHI and BspMII. The linker fragment is then ligated to the 3' end of the LIF-R fragment (cleaved at its 3' terminus after the Gly₄Ser sequence with BamHI). The ligation is conducted under conventional conditions. The 3' end of the

WO 93/10151

PCT/US92/10272

35

linker fragment is ligated to the BspMII-cleaved 5' end of the gp130 fragment. The resulting DNA fragment encodes a receptor of the present invention comprising (from 5' to 3') the signal sequence and extracellular domain of LIF-R attached to a (Gly₄Ser)₈ polypeptide linker which is attached to the mature coding sequence of the gp130 extracellular domain.

This DNA fragment may be inserted into any suitable cloning and/or expression vector. For example, the pBLUESCRIPT SK[®] vector may be digested with SalI and NotI and the ligated DNA fragment inserted therein. *E. coli* cells are then transformed with the recombinant vector by conventional procedures.

In an alternative procedure, the pBLUESCRIPT SK[®] vector containing the Ser(Gly₄Ser)₆Gly linker sequence is digested with SalI and BamHI and the above described LIF-R-encoding fragment is inserted therein. The resulting vector is then digested with BspMII and NotI and the gp130-encoding fragment is then inserted to form the DNA sequence encoding the receptor of the present invention. The cloned receptor-encoding DNA fragment may be excised and inserted into any suitable expression vector (chosen according to the type of host cell that is desired) using conventional procedures. Host cells transformed with the recombinant expression vector are cultivated to produce the receptor protein. Mammalian host cells are generally preferred for producing the recombinant receptor fusion proteins of the present invention.

The receptor-encoding construct may be excised by SalI and NotI digestion and inserted into a vector suitable for use in mammalian host cells. One suitable vector is designated pDC406. cDNA molecules inserted at the SalI site of this vector are transcribed and translated using regulatory elements derived from HIV and adenovirus. pDC406 contains origins of replication derived from SV40, Epstein-Barr virus and pBR322. The pDC406 vector into which interleukin-1 receptor cDNA has been cloned has been deposited with the American Type Culture Collection, Rockville, Maryland USA under accession number CRL10478. The interleukin-1 receptor cDNA may be excised from the vector using conventional techniques and replaced with the receptor encoding DNA of the present invention prepared above. pDC406 is a derivative of HAV-EO described by Dower et al., *J. Immunol.* 142:4314 (1989). pDC406 differs from HAV-EO by the deletion of the intron present in the adenovirus 2 tripartite leader sequence in HAV-EO.

Examples of suitable mammalian cells for expressing a receptor fusion protein include CV-1 cells (ATCC CCL70) and COS-7 cells, (ATCC CRL 1651) both derived from monkey kidney. Another monkey kidney cell line CV-1/EBNA (ATCC CRL 10478) was derived by transfection of the CV-1 cell line with a gene encoding Epstein-

WO 93/10151

PCT/US92/10272

36

Barr virus nuclear antigen-1 (EBNA-1) and with a vector containing CMV regulatory sequences. See McMahan et al., *EMBO J.* 10:2821 (1991). The EBNA-1 gene allows for episomal replication of expression vectors, such as HAV-EO or pDC406, that contain the EBV origin of replication.

5

Example 4

Preparation of a Recombinant Receptor Fusion Protein Designating gp130-Linker-LIF-R

10 This receptor of the present invention differs from that of Example 3 in that the LIF-R polypeptide (which was the 5' polypeptide in the receptor of Example 3) is now the 3' polypeptide. The following oligonucleotides were synthesized for use in preparing the fusion protein:

15 SEQ ID NO: 12

5' GATATGTCGACAAGATGTTGACGTTGCAGACTTGG 3' (oligonucleotide no. 5)

SEQ ID NO: 13

3' CAAACGAGTTCCTCTTTAACTTCCTCCTCCACCTAGGTACG 5' (oligonucleotide no. 6)

20

SEQ ID NO: 14

5' CGCGTCCGGAGGAGGTGGTAGCCAGAAAAAGGGGGCTCCTCATG 3' (oligonucleotide
no. 7)

25 SEQ ID NO: 15

3' CATACATACACCACTGTTTCCTTTTAAAGATCGCCGGCGTACG 5' (oligonucleotide no.
8)

30 Oligonucleotides 5 and 6 are employed in a polymerase chain reaction procedure to isolate a fragment of gp130. The 5' primer (oligonucleotide number 5) includes nucleotides 244 to 264 of SEQ ID NO: 1, (the sequence encoding the first seven amino acids of the gp130 signal sequence). Oligonucleotide number 5 also includes a sequence that introduces an upstream SalI site. This nucleotide sequence is capable of annealing to the (-) strand that is complementary to nucleotides 244 to 264 of SEQ ID
35 NO: 1. The 3' primer (oligonucleotide number 6) includes a sequence complementary to nucleotides 2080 to 2100 of SEQ ID NO: 1, i.e. includes antisense nucleotides encoding the last seven amino acids of the gp-130 extracellular domain.

WO 93/10151

PCT/US92/10272

37

Oligonucleotide number 6 also encodes a Gly₄Ser sequence immediately 3' to (and in phase with) the gp130 sequence, and also inserts a downstream BamHI site.

5 A PCR reaction is conducted as described in Example 3 but employing oligonucleotides 5 and 6 on the gp130 cDNA template. A DNA sequence encoding a gp130 fragment that includes the 5' signal sequence and the entire extracellular domain, but none of the transmembrane region or the cytoplasmic domain, is isolated by the PCR reaction. A Gly₄Ser sequence is fused to the 3' terminus of the gp130 fragment. The PCR reaction products are digested with SalI and BamHI and the desired fragment is isolated.

10 An LIF-R fragment is isolated and amplified by a PCR reaction employing oligonucleotides 7 and 8. The 5' primer (oligonucleotide number 7) includes nucleotides 311 to 331 of SEQ ID NO: 5, which encode the first seven amino acids of the mature LIF-R protein. This nucleotide sequence is capable of annealing to the (-) strand complementary to nucleotides 311 to 331 of SEQ ID NO: 5. Oligonucleotide
15 number 7 also encodes a Gly₄Ser sequence fused to the 5' end of the LIF-R sequence, and inserts an upstream BspMII site. The 3' primer (oligonucleotide number 8) is complementary to nucleotides 2651 to 2677 of SEQ ID NO: 5 (which encode the last nine amino acids of the LIF-R extracellular domain.) Oligonucleotide number 8 also adds a stop codon at the 3' end of the LIF-R sequence, and inserts a Not I site
20 downstream. The PCR reaction products are digested with BspMII and NotI and the desired fragment is isolated.

A DNA sequence encoding the desired receptor protein is prepared by ligating the BamHI site of the gp130 fragment prepared above to the BamHI site at the 5' terminus of the linker fragment described in Example 3. Likewise the C-terminus of
25 the linker encoding fragment is ligated at the BspMII site to the complementary site of the LIF-R encoding fragment prepared above. The resulting DNA fragment may be cloned into an expression vector using procedures described in Example 3. The receptor encoded by the isolated DNA fragment comprises (from the N-terminus to the C-terminus) the signal sequence and extracellular domain of gp130 attached to a
30 (Gly₄Ser)₈ polypeptide linker which is attached to the mature coding sequence of the LIF-R extracellular domain.

PCI/US92/10272

WO 93/10151

38

Example 5

Receptor Fusion Protein Comprising LIF-R Attached to gp-130 Through An Fc Polypeptide Linker

5

A receptor prepared in accordance with the following procedures is depicted in Figure 4. The following oligonucleotides were synthesized for use in preparing the receptor fusion protein:

SEQ ID NO: 16

10

3' CATACATACACCACTGTTTCCTTTTAAGACTCGGGTCTAGATACG 5' (oligonucleotide no. 9)

SEQ ID NO: 17

3' CAAACGAGTTTCCTCTTAACTTCTCGGGTCTAGATACG 5' (oligonucleotide no. 10)

15

An LIF-R encoding DNA sequence is isolated and amplified in a PCR reaction using oligonucleotides 1 and 9. Oligonucleotide number 1 (the 5' primer) inserts an upstream SalI site and has been described in Example 3. The 3' primer is oligonucleotide number 9 which includes a sequence complementary to nucleotides 2651 to 2677 of SEQ ID NO: 5, i.e., includes antisense nucleotides encoding the last nine amino acids of the extracellular domain of LIF-R. Oligonucleotide number 9 also inserts a downstream BglII site. The PCR reaction products are digested with SalI and BglII, and the desired LIF-R encoding DNA fragment is isolated by gel electrophoresis using conventional procedures. Due to the presence of an internal BglII site in the LIF-R sequence, the BglII digestion should be carried out under conditions that effect partial digestion.

25

A gp130 encoding DNA fragment is isolated and amplified by PCR reaction using oligonucleotides 5 and 10. The 5' primer (oligonucleotide number 5) inserts an upstream SalI site and has been described above in Example 4. The 3' primer is oligonucleotide number 10, which includes a sequence complementary to nucleotides 2080 to 2100 of SEQ ID NO: 1, i.e., includes antisense nucleotides encoding the last seven amino acids of the gp130 extracellular domain. Oligonucleotide number 10 also inserts a downstream BglII site. The PCR reaction products are digested with SalI and BglII, and the desired gp130 encoding DNA fragment is isolated by gel electrophoresis using conventional techniques.

35

cDNA encoding a single chain polypeptide derived from the Fc region of a human IgG1 antibody has been cloned into the above-described pBLUESCRIPT SK®

WO 93/10151

PCT/US92/10272

39

vector to produce a recombinant vector designated hlgGIFc. A unique BglII site is positioned near the 5' end of the inserted Fc encoding sequence. An SpeI site is immediately downstream of the stop codon. The DNA and encoded amino acid sequences of the cloned Fc cDNA are presented in SEQ ID NO: 3 and SEQ ID NO: 4.

5 The Fc polypeptide encoded by the cDNA extends from the N-terminal hinge region to the native C-terminus, i.e., is an essentially full-length antibody Fc region. Fc fragments, e.g., those that are truncated at the C-terminal end, also may be employed. The fragments should contain multiple cysteine residues (at least the cysteine residues in the hinge region). The antibody from which the Fc polypeptide is
10 derived is preferably of the same species as the patient to be treated with the fusion protein prepared therefrom.

Plasmid hlgGIFc is digested with BglII and SalI and the BglII/SalI LIF-R fragment prepared above is ligated into the vector by conventional techniques. The Fc
15 encoding sequence is positioned downstream of, and in the same reading frame as, the LIF-R sequence. In a separate reaction, the above-described SalI/BglII fragment of gp130 is also inserted into the same vector. Plasmid vectors containing the desired DNA insert are identified by restriction endonuclease digestion analysis, using
convention techniques.

The cloned DNA segment encoding the LIF-R-Fc fusion polypeptide may be
20 excised from the pBLUESCRIPT SK[®] vector by digestion with SalI and NotI. Likewise, the DNA segment encoding the gp130-Fc fusion polypeptide may be excised by SalI/NotI digestion. Each excised DNA segment is inserted into an appropriate expression vector, depending on the type of host cell that is desired. One suitable
expression vector is the plasmid pDC406, which may be transformed into mammalian
25 host cells as described in Example 3.

In one embodiment of the invention, an expression vector encoding the LIF-R-Fc fusion and a second expression vector encoding the gp130-Fc fusion are co-
transfected into the desired host cells. Two separate recombinant polypeptides are thus
produced in the host cells. The first polypeptide comprises the Fc polypeptide fused in
30 frame to the C-terminus of the gp130 fragment. The second polypeptide comprises the Fc polypeptide fused in frame to the C-terminus of the LIF-R fragment. Disulfide
bonds that form between the two Fc regions covalently link the two separate fusion
polypeptides into a receptor protein of the present invention.

Alternatively, the LIF-R-Fc and gp130-Fc polypeptides may be separately
35 transformed into host cells (as opposed to co-transfection into the same host cell.) The two polypeptides are purified from the host cells and then combined in a suitable

WO 93/10151

PCI/US92/10272

40

buffered solution, whereupon interchain disulfide bonds form between the two Fc regions.

5 The receptor protein may be purified using any of a number of conventional protein purification techniques. Since antibody Fc regions bind to protein A and protein G, affinity chromatography employing protein A or protein G attached to an insoluble support material may be employed in the purification process. In one procedure, one liter of culture supernatant containing the receptor is passed over a solid phase protein G column, and the column is then washed thoroughly with phosphate-buffered saline (PBS). The adsorbed Fc-containing fusion protein is eluted with 50 mM glycine buffer, pH 3 and brought to pH 7 with 2 M Tris buffer, pH 9. Further purification may involve immunoaffinity column(s), e.g., affinity columns having LIF or OSM bound thereto.

Example 6

15 Preparation of Monoclonal Antibodies Directed against a Receptor

Preparations of a purified receptor protein of the present invention, or transfected COS cells expressing high levels of the receptor, are employed to generate monoclonal antibodies against the receptor using conventional techniques, for example, those disclosed in U.S. Patent 4,411,993. To immunize mice, a receptor immunogen is emulsified in complete Freund's adjuvant and injected subcutaneously in amounts ranging from 10-100µg into Balb/c mice. Ten to twelve days later, the immunized animals are boosted with additional immunogen emulsified in incomplete Freund's adjuvant and periodically boosted thereafter on a weekly to biweekly immunization schedule. Serum samples are periodically taken by retro-orbital bleeding or tail-tip excision for testing by dot-blot assay (antibody sandwich) or ELISA (enzyme-linked immunosorbent assay). Other assay procedures are also suitable. Following detection of an appropriate antibody titer, positive animals are given an intravenous injection of antigen in saline. Three to four days later, the animals are sacrificed, splenocytes harvested, and fused to the murine myeloma cell line NS1. Hybridoma cell lines generated by this procedure are plated in multiple microtiter plates in a HAT selective medium (hypoxanthine, aminopterin, and thymidine) to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

35 Hybridoma clones thus generated can be screened by ELISA for reactivity with the receptor protein, for example, by adaptations of the techniques disclosed by Engvall et al., *Immunochem* 8:871 (1971) and in U.S. Patent 4,704,004. Positive clones are then injected into the peritoneal cavities of syngeneic Balb/c mice to produce ascites

WO 93/10151

PCT/US92/10272

41

containing high concentrations (greater than 1 mg/ml) of anti-receptor monoclonal antibody. The resulting monoclonal antibody can be purified by ammonium sulfate precipitation followed by gel exclusion chromatography, and/or affinity chromatography based on binding of antibody to Protein A of *Staphylococcus aureus*.

5

Example 7

Heterodimeric Receptor

10 An expression vector encoding a fragment of the human LIF-R extracellular domain fused to a polypeptide derived from the Fc region of an antibody was constructed as follows. A second expression vector encoding a fragment of the human gp130 extracellular domain fused to an Fc polypeptide also was constructed.

15 Plasmid pHLIF-R-65 (ATCC 68491), which contains human LIF-R cDNA in expression vector pDC303 as described in example 3, was digested with the restriction enzymes *Asp718* and *XmnI*. *Asp718* cleaves the vector upstream of the LIF-R cDNA insert. *XmnI* is a blunt cutter that cleaves within the codon for amino acid number 702 (Asp) of SEQ ID NO:5, upstream of the transmembrane region. The desired *Asp718/XmnI* fragment (about 2,444 bp in length) was separated by electrophoresis on an agarose gel and purified by conventional procedures using an Elutip column.

20 A recombinant vector designated hIgG1Fc, comprising cDNA encoding a single chain polypeptide derived from the Fc region of a human IgG1 antibody in a pBLUESCRIPT SK® vector was described in example 5. The DNA and encoded amino acid sequences of the cloned Fc cDNA are presented in SEQ ID NO:3 and SEQ ID NO:4. A polylinker region comprising a number of restriction sites is positioned
25 immediately upstream of the Fc cDNA.

Plasmid hIgG1Fc was digested with *Asp718* and *SnaI*, which cleave within the polylinker upstream of the Fc sequence. The *Asp718/XmnI* LIF-R fragment prepared above was ligated into the cleaved hIgG1Fc vector by conventional techniques. *SnaI* and *XmnI* both produce blunt ends, which will ligate together. In the resulting
30 recombinant vector, the Fc encoding sequence is positioned downstream of, and in the same reading frame as, the LIF-R sequence. The encoded LIF-R/Fc fusion protein comprises amino acids -44 to 702 of SEQ ID NO:5 (LIF-R), followed by six amino acids constituting a peptide linker encoded by the polylinker segment of plasmid hIgG1Fc, followed by amino acids 1-232 of SEQ ID NO:3 (Fc). *E. coli* cells were
35 transformed with the ligation mixture and plasmids were isolated therefrom by standard procedures. Plasmid vectors containing the desired DNA insert were identified by restriction endonuclease digestion analysis.

WO 93/10151

PCI/US92/10272

42

The cloned DNA segment encoding the LIF-R/Fc fusion polypeptide was excised from the recombinant vector by digestion with *Asp718* and *NotI*. The *NotI* enzyme cleaves the vector in a polylinker region just downstream of the Fc cDNA insert. The excised DNA segment (3.2 kb) is inserted into an appropriate expression vector, depending on the type of host cell that is desired. One suitable expression vector is pCAV/NOT, a mammalian expression vector described in PCT application WO 90/05183.

pCAV/NOT was cleaved with *Asp718* and *NotI*, both of which cleave in the multiple cloning site. The LIF-R/Fc-encoding *Asp718/NotI* DNA fragment prepared above was ligated into the vector.

An expression vector encoding a soluble gp130/Fc fusion protein was constructed as follows. Recombinant vector B10G/pDC303 (ATCC 68827) comprising human gp130 cDNA (described in example 3) was digested with *EcoRI*, and the resulting 5' overhang was rendered blunt using T4 DNA polymerase. The recognition site for *EcoRI* comprises nucleotides 2056-2061 of SEQ ID NO:1. The *EcoRI*-digested vector was then cleaved with *XhoI*, which cleaves in the vector upstream of the gp130 cDNA insert.

Plasmid hIgG1Fc, comprising Fc polypeptide-encoding cDNA as described above, was digested with *SnaI* (a blunt cutter) and *NotI*, which cleave upstream and downstream, respectively, of the inserted Fc cDNA. The (*EcoRI*)/*XhoI* gp130 fragment isolated above was ligated to the Fc-containing fragment and to *XhoI/NotI*-digested vector SF CAV/NOT. The mammalian expression vector SF CAV/NOT is essentially identical to SF CAV (ATCC 68922) but contains a *NotI* site. SF CAV/NOT also is essentially identical to pCAV/NOT, described in PCT application WO 90/05183, except that a segment of the adenovirus-2 tripartite leader (TPL) containing a cryptic promoter functional in bacteria has been deleted. Protein expression from the cryptic promoter is potentially disadvantageous for preparing and isolating a desired recombinant plasmid in bacterial cells.

E. coli cells were transformed with the ligation mixture, plasmids were isolated therefrom by conventional procedures, and the desired recombinant plasmids were identified by restriction analysis. The gp130/Fc fusion protein encoded by the desired recombinant vector comprises (from N- to C-terminus) amino acids -22 to 582 of SEQ ID NO:2 (gp130), followed by 7 amino acids constituting a peptide linker encoded by the polylinker segment of plasmid hIgG1Fc, followed by amino acids 1-232 of SEQ ID NO:4 (Fc).

COS-7 cells (ATCC CRL 1651) were transfected with either the LIF-R/Fc-encoding recombinant expression vector or the gp130/Fc-encoding expression vector

WO 93/10151

PCT/US92/10272

43

prepared above, or with both expression vectors. The cells were cultivated to allow expression of the soluble fusion proteins. The expressed proteins were recovered by incubating culture supernatant with Protein G Sepharose beads (available from Pharmacia) overnight at 4°C, then pelleting the beads by centrifugation. The binding of ¹²⁵I-labeled human oncostatin M and ¹²⁵I-labeled human LIF by the proteins bound to the beads was analyzed.

Binding affinity was determined by performing a variation of a standard Scatchard analysis. The binding assay procedure was similar to that described by Mosley et al. (*Cell* 59:335, 1989) except that the proteins, being soluble, are attached to Protein G Sepharose beads rather than being on the surface of the transfected cells. Briefly, in a 96-well microtiter plate, each of ten 1:2 serial dilutions of ¹²⁵I-LIF or ¹²⁵I-oncostatin M was incubated with a sample comprising the expressed proteins (bound to the beads) resuspended in RPMI 1640 containing 2.5% bovine serum albumin, 0.2% (v/v) sodium azide and 20mM Hepes, pH 7.4, for 2 hours at 4°C with agitation. Duplicate standard cold competition wells also were incubated. Centrifuge tubes containing bovine calf serum were used in place of the phthalate oil mixture-containing tubes in the separation method described by Dower et al., *J. Immunol.* 132:751 (1984) and Park et al., *J. Biol. Chem.* 261:4177 (1986) and in example 1 above. Aliquots of each incubation mixture were transferred to the tubes. After centrifugation, tubes were cut, the radioactivity counted, and processed as for standard Scatchard analysis.

Figure 7 presents Scatchard analyses of the binding of ¹²⁵I oncostatin M by gp130/Fc homodimers produced by the cells transfected with the gp130/Fc vector alone (upper left) and by the proteins expressed by the co-transfected cells (lower left). Scatchard analyses of the binding of ¹²⁵I LIF by LIF-R/Fc homodimers produced by the cells transfected with the LIF-R/Fc vector alone (upper right) and by the proteins expressed by the co-transfected cells (lower right) are also presented in figure 7. A shift toward higher affinity binding of oncostatin M by the proteins recovered from the co-transfected cells, compared to the gp130/Fc homodimer, is evident from figure 7. Likewise, the data in figure 7 indicate a shift toward higher affinity binding of LIF by the proteins recovered from the co-transfected cells, compared to the LIF-R/Fc homodimer. The shift toward higher affinity binding indicates the presence of heterodimers comprising LIF-R/Fc and gp130/Fc, and further indicates that the LIF-R and gp130 moieties are cooperating, i.e., interacting, in the binding of oncostatin M and LIF. Controls demonstrated no oncostatin M binding by LIF-R homodimers, and no LIF binding by gp130 homodimers.

WO 93/10151

PCT/US92/10272

44

Example 8

Receptors Comprising LIF-R and gp130 Polypeptides Lacking FNIII Domains

5 DNA sequences encoding soluble LIF-R and gp130 proteins lacking fibronectin type III (FNIII) domains were isolated and fused to an Fc-encoding sequence. Deleting the FNIII domains affords the advantage of reducing the size of the LIF-R/Fc and gp130/Fc fusion proteins. The LIF-R protein of SEQ ID NO:6 comprises three repeats of a fibronectin type III-like module in the extracellular domain. The three domains containing FNIII modules comprise amino acids 487 (Thr) to 584 (Asn), 585 (Asp) to 679 (Ala), and 680 (Pro) to 789 (Ser), respectively, of SEQ ID NO:6. gp130 also 10 contains three FNIII domains, comprising amino acids 300 (Tyr) to 399 (Phe), 400 (Gln) to 496 (Pro), and 497 (Pro) to 597 (Glu), respectively, of SEQ ID NO:2. From one to all three of the FNIII domains may be removed from gp130 or LIF-R to reduce the size of the protein.

15 The FNIII domains of human LIF-R were removed by digesting the LIF-R/Fc-encoding expression vector prepared in example 7 with the restriction endonuclease Eco O 109I (isoschizomer of Dra II) and filling in the resulting overhangs using T4 DNA polymerase according to conventional procedures. The recognition site for Eco O 109I spans nucleotides 1789-1795 of SEQ ID NO:5 (LIF-R), cleaving within the codons for amino acids 8-9 of the first FNIII domain of LIF-R. The cleaved vector was then 20 digested with BstXI and EcoR5. The recognition site for BstXI spans nucleotides 1048-1059 of SEQ ID NO:5 and EcoR5 (which generates blunt ends) cleaves within the polylinker upstream of the Fc sequence. The BstXI/EcoR5 fragment (comprising the 5' end of LIF-R, the vector, the entire Fc sequence, and a portion of the polylinker) 25 and the BstXI/(Eco O 109I) LIF-R fragment were isolated and ligated together. *E. coli* cells were transformed with the ligation mixture, plasmids were isolated therefrom, and the desired recombinant plasmid was identified by restriction analysis. The resulting construct encodes a fusion protein comprising (from N- to C-terminus) amino acids -44 to 494 of SEQ ID NO:5 (LIF-R), a four amino acid spacer peptide -His-Arg-Tyr-Val- encoded by the polylinker segment, and amino acids 1-232 of SEQ ID NO:3 (Fc). The 30 LIF-R polypeptide moiety contains the first 8 amino acids of the first FNIII domain, but lacks the remainder of the first FNIII domain and all of the second and third FNIII domains.

35 The FNIII domains of gp130 were removed by digesting the recombinant gp130/Fc-encoding expression vector prepared in example 7 with BstXI, then blunting the overhang using T4 DNA polymerase according to conventional procedures. The recognition site for BstXI spans nucleotides 1231-1242 of SEQ ID NO:1 (gp130).

WO 93/10151

PCT/US92/10272

45

cleaving within the codons for amino acids 10-11 of the first FNIII domain of gp130. The cleaved vector was then digested with EcoR5, which cleaves within the polylinker upstream of the Fc sequence and generates blunt ends. The (BstX1)/EcoR5 fragment comprising the 5' end of gp130 (lacking the FNIII domains), the vector sequence, the Fc sequence, and a portion of the polylinker, was ligated. *E. coli* cells were transformed with the ligation mixture, plasmids were isolated therefrom, and the desired recombinant plasmid was identified by restriction analysis. The fusion protein encoded by the construct comprises (from N- to C-terminus) amino acids -22 to 308 of SEQ ID NO:2 (gp130), a four amino acid spacer peptide -Asn-Arg-Tyr-Val- encoded by the polylinker segment, and amino acids 1-232 of SEQ ID NO:3 (Fc). The gp130 polypeptide moiety contains the first 9 amino acids of the first FNIII domain, but lacks the remainder of the first FNIII domain and all of the second and third FNIII domains.

FNIII domains may be deleted from the gp130 component of a receptor of the present invention, from the LIF-R component, or from both. In one embodiment of the invention, COS-7 cells were co-transfected with the soluble LIF-R/Fc-encoding mammalian expression vector prepared in example 7 and the mammalian expression vector encoding a soluble gp130/Fc protein lacking the FNIII domains prepared above. Analysis of the expressed proteins by SDS-PAGE revealed a band of the molecular weight expected for the heterodimer, along with bands that include those of the molecular weight expected for the two homodimers. Scatchard analyses conducted according to the procedures described in example 7 demonstrated a shift toward higher affinity binding of LIF and oncostatin M for proteins expressed by the co-transfected cells compared to the corresponding homodimers. This result indicates the presence of heterodimers comprising LIF-R/Fc and gp130/Fc, and further indicates that the LIF-R and gp130 moieties are cooperating, i.e., interacting, in binding oncostatin M and LIF.

BRIEF DESCRIPTION OF THE SEQUENCE LISTING

SEQ ID NO:1 and SEQ ID NO:2 present the DNA sequence and encoded amino acid sequence for cloned cDNA encoding an N-terminal fragment of gp130.

SEQ ID NO:3 and SEQ ID NO:4 present the DNA sequence and encoded amino acid sequence for cloned cDNA encoding a polypeptide that corresponds to the Fc region of an IgG1 antibody.

SEQ ID NO:5 and SEQ ID NO:6 present the DNA sequence and encoded amino acid sequence for cloned cDNA encoding an N-terminal fragment of LIF-R.

WO 93/10151

PCT/US92/10272

46

SEQ ID NO:7 presents the DNA sequence of the coding strand of a chemically synthesized DNA molecule encoding a polypeptide linker used in constructing certain receptors of the present invention.

5

SEQ ID NO:8 - SEQ ID NO:17 present the DNA sequence of various single-stranded oligonucleotide primers employed in polymerase chain reactions to construct certain receptors of the present invention.

WO 93/10151

PCT/US92/10272

47
SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Gearing, David P.

(ii) TITLE OF INVENTION: Receptor for Oncostatin M and Leukemia
Inhibitory Factor

(iii) NUMBER OF SEQUENCES: 17

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Immunex Corporation
(B) STREET: 51 University Street
(C) CITY: Seattle
(D) STATE: WA
(E) COUNTRY: USA
(F) ZIP: 98101

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.23

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Seese, Kathryn A.
(B) REGISTRATION NUMBER: 32,172
(C) REFERENCE/DOCKET NUMBER: 2607

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 206-587-0430
(B) TELEFAX: 206-587-0606

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2369 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: N-terminal

(vi) ORIGINAL SOURCE:

(F) TISSUE TYPE: human placenta

(vii) IMMEDIATE SOURCE:

PCT/US92/10272

WO 93/10151

48

(B) CLONE: B10G/pDC303

(ix) FEATURE:

(A) NAME/KEY: CD8

(B) LOCATION: 244..2369

(ix) FEATURE:

(A) NAME/KEY: mat_peptide

(B) LOCATION: 310..2369

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: 244..309

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GGCCCCGCGGA GTCCGCTGG GCGCCCCCG CCGAGCTGAA CCGGGGGCCG CGCCTGCCAG	60
GCCGACGGGT CTGGCCGAGC CTGGCGCCAA GGGGTTCGTG CGCTGTGGAG ACGCGGAGGG	120
TCGAGGCGGC GCGGCCTGAG TGAAACCCAA TGGAAAAGC ATGACATTTA GAAGTAGAAG	180
ACTTAGCITC AAATCCCTAC TCCTTCACTT ACTAATTTTG TGATTGGAA ATATCCGCGC	240
AAG ATG TTG ACG TTG CAG ACT TGG CTA GTG CAA GCC TTG TTT ATT TTC	288
Met Leu Thr Leu Gln Thr Trp Leu Val Gln Ala Leu Phe Ile Phe	
-22 -20 -15 -10	
CTC ACC ACT GAA TCT ACA GGT GAA CTT CTA GAT CCA TGT GGT TAT ATC	336
Leu Thr Thr Glu Ser Thr Gly Glu Leu Leu Asp Pro Cys Gly Tyr Ile	
-5 1 5	
AGT CCT GAA ICT CCA GTT GTA CAA CTT CAT TCT AAT TTC ACT GCA GTT	384
Ser Pro Glu Ser Pro Val Val Gln Leu His Ser Asn Phe Thr Ala Val	
10 15 20 25	
TGT GTG CTA AAG GAA AAA TGT ATG GAT TAT TTT CAT GTA AAT GCT AAT	432
Cys Val Leu Lys Gln Lys Cys Met Asp Tyr Phe His Val Asn Ala Asn	
30 35 40	
TAC ATT GTC TGG AAA ACA AAC CAT TTT ACT ATT CCT AAG GAG CAA TAT	480
Tyr Ile Val Trp Lys Thr Asn His Phe Thr Ile Pro Lys Glu Gln Tyr	
45 50 55	
ACT ATC ATA AAC AGA ACA GCA TCC AGT GTC ACC TTT ACA GAT ATA GCT	528
Thr Ile Ile Asn Arg Thr Ala Ser Ser Val Thr Phe Thr Asp Ile Ala	
60 65 70	
TCA TTA AAT ATT CAG CTC ACT TGC AAC ATT CTT ACA TTC GGA CAG CTT	576
Ser Leu Asn Ile Gln Leu Thr Cys Asn Ile Leu Thr Phe Gly Gln Leu	
75 80 85	
GAA CAG AAT GTT TAT GGA ATC ACA ATA ATT TCA GGC TTG CCT CCA GAA	624
Gln Gln Asn Val Tyr Gly Ile Thr Ile Ile Ser Gly Leu Pro Pro Glu	
90 95 100 105	
AAA CCT AAA AAT TTG AGT TGC ATT GTG AAC GAG GGG AAG AAA ATG AGG	672
Lys Pro Lys Asn Leu Ser Cys Ile Val Asn Glu Gly Lys Lys Met Arg	
110 115 120	

WO 93/10151

PCT/US92/10272

49																	
TGT	CAG	TGG	GAT	GGT	GGA	AGG	GAA	ACA	CAC	TTG	GAG	ACA	AAC	TTC	ACT		720
Cys	Glu	Trp	Asp	Gly	Gly	Arg	Glu	Thr	His	Leu	Glu	Thr	Asn	Phe	Thr		
			125					130					135				
TTA	AAA	TCT	GAA	TGG	GCA	ACA	CAC	AAG	TTT	GCT	GAT	TGC	AAA	GCA	AAA		768
Leu	Lys	Ser	Glu	Trp	Ala	Thr	His	Lys	Phe	Ala	Asp	Cys	Lys	Ala	Lys		
		140					145					150					
CGT	GAC	ACC	CCC	ACC	TCA	TGC	ACT	GTT	GAT	TAT	TCT	ACT	GTG	TAT	TTT		816
Arg	Asp	Thr	Pro	Thr	Ser	Cys	Thr	Val	Asp	Tyr	Ser	Thr	Val	Tyr	Phe		
	155					160					165						
GTC	AAC	ATT	GAA	GTC	TGG	GTA	GAA	GCA	GAG	AAT	GCC	CTT	GGG	AAG	GTT		864
Val	Asn	Ile	Glu	Val	Trp	Val	Glu	Ala	Glu	Asn	Ala	Leu	Gly	Lys	Val		
170					175					180					185		
ACA	TCA	GAT	CAT	ATC	AAT	TTT	GAT	CCT	GTA	TAT	AAA	GTG	AAG	CCC	AAT		912
Thr	Ser	Asp	His	Ile	Asn	Phe	Asp	Pro	Val	Tyr	Lys	Val	Lys	Pro	Asn		
			190						195					200			
CCG	CCA	CAT	AAT	TTA	TCA	GTG	ATC	AAC	TCA	GAG	GAA	CTG	TCT	AGT	ATC		960
Pro	Pro	His	Asn	Leu	Ser	Val	Ile	Asn	Ser	Glu	Glu	Leu	Ser	Ser	Ile		
			205					210					215				
TTA	AAA	TTG	ACA	TGG	ACC	AAC	CCA	AGT	ATT	AAG	AGT	GTT	ATA	ATA	CTA		1008
Leu	Lys	Leu	Thr	Trp	Thr	Asn	Pro	Ser	Ile	Lys	Ser	Val	Ile	Ile	Leu		
		220					225					230					
AAA	TAT	AAC	ATT	CAA	TAT	AGG	ACC	AAA	GAT	GCC	TCA	ACT	TGG	AGC	CAG		1056
Lys	Tyr	Asn	Ile	Gln	Tyr	Arg	Thr	Lys	Asp	Ala	Ser	Thr	Trp	Ser	Gln		
	235					240				245							
ATT	CCT	CCT	GAA	GAC	ACA	GCA	TCC	ACC	CGA	TCT	TCA	TTC	ACT	GTC	CAA		1104
Ile	Pro	Pro	Glu	Asp	Thr	Ala	Ser	Thr	Arg	Ser	Ser	Phe	Thr	Val	Gln		
250					255					260					265		
GAC	CTT	AAA	CCT	TTT	ACA	GAA	TAT	GTG	TTT	AGG	ATT	CGC	TGT	ATG	AAG		1152
Asp	Leu	Lys	Pro	Phe	Thr	Glu	Tyr	Val	Phe	Arg	Ile	Arg	Cys	Met	Lys		
				270					275					280			
GAA	GAT	GGT	AAG	GGA	TAC	TGG	AGT	GAC	TGG	AGT	GAA	GAA	GCA	AGT	GGG		1200
Glu	Asp	Gly	Lys	Gly	Tyr	Trp	Ser	Asp	Trp	Ser	Glu	Glu	Ala	Ser	Gly		
			285					290					295				
ATC	ACC	TAT	GAA	GAT	AGA	CCA	TCT	AAA	GCA	CCA	AGT	TTC	TGG	TAT	AAA		1248
Ile	Thr	Tyr	Glu	Asp	Arg	Pro	Ser	Lys	Ala	Pro	Ser	Phe	Trp	Tyr	Lys		
		300					305					310					
ATA	GAT	CCA	TCC	CAT	ACT	CAA	GGC	TAC	AGA	ACT	GTA	CAA	CTC	GTG	TGG		1296
Ile	Asp	Pro	Ser	His	Thr	Gln	Gly	Tyr	Arg	Thr	Val	Gln	Leu	Val	Trp		
	315					320					325						
AAG	ACA	TTG	CCT	CCT	TTT	GAA	GCC	AAT	GGA	AAA	ATC	TTG	GAT	TAT	GAA		1344
Lys	Thr	Leu	Pro	Pro	Phe	Glu	Ala	Asn	Gly	Lys	Ile	Leu	Asp	Tyr	Glu		
330					335					340					345		
GTG	ACT	CTC	ACA	AGA	TGG	AAA	TCA	CAT	TTA	CAA	AAT	TAC	ACA	GTT	AAT		1392
Val	Thr	Leu	Thr	Arg	Trp	Lys	Ser	His	Leu	Gln	Asn	Tyr	Thr	Val	Asn		
				350					355					360			

30															1440	
GCC	ACA	AAA	CTG	ACA	GTA	AAT	CTC	ACA	AAT	GAT	CGC	TAT	CTA	GCA	ACC	1440
Ala	Thr	Lys	Leu	Thr	Val	Asn	Leu	Thr	Asn	Asp	Arg	Tyr	Leu	Ala	Thr	
365															375	
370																
1488																
CTA	ACA	GTA	AGA	AAT	CTT	GTT	GGC	AAA	TCA	GAT	GCA	GCT	GTT	TTA	ACT	1488
Leu	Thr	Val	Arg	Asn	Leu	Val	Gly	Lys	Ser	Asp	Ala	Ala	Val	Leu	Thr	
380															390	
385																
1536																
ATC	CCT	GCC	TGT	GAC	TTT	CAA	GCT	ACT	CAC	CCT	GTA	ATG	GAT	CTT	AAA	1536
Ile	Pro	Ala	Cys	Asp	Phe	Gln	Ala	Thr	His	Pro	Val	Met	Asp	Leu	Lys	
395															405	
400																
1584																
GCA	TTC	CCC	AAA	GAT	AAC	ATG	CTT	TGG	GTG	GAA	TGG	ACT	ACT	CCA	AGG	1584
Ala	Phe	Pro	Lys	Asp	Asn	Met	Leu	Trp	Val	Glu	Trp	Thr	Thr	Pro	Arg	
410															425	
415															420	
1632																
GAA	TCT	GTA	BAQ	AAA	TAT	ATA	CTT	GAG	TGG	TGT	GTG	TTA	TCA	GAT	AAA	1632
Glu	Ser	Val	Lys	Lys	Tyr	Ile	Leu	Glu	Trp	Cys	Val	Leu	Ser	Asp	Lys	
430															440	
435																
1680																
GCA	CCC	TGT	ATC	ACA	GAC	TGG	CAA	CRA	GAA	GAT	GGT	ACC	GTG	CAT	CGC	1680
Ala	Pro	Cys	Ile	Thr	Asp	Trp	Gln	Gln	Glu	Asp	Gly	Thr	Val	His	Arg	
445															455	
450																
1728																
ACC	TAT	TTA	AGA	GGG	AAC	TTA	GCA	GAG	AGC	AAA	TGC	TAT	TTG	ATA	ACA	1728
Thr	Tyr	Leu	Arg	Gly	Asn	Leu	Ala	Glu	Ser	Lys	Cys	Tyr	Leu	Ile	Thr	
460															470	
465																
1776																
GTT	ACT	CCA	GTA	TAT	GCT	GAT	GGA	CCA	GGA	AGC	CCT	GAA	TCC	ATA	AAG	1776
Val	Thr	Pro	Val	Tyr	Ala	Asp	Gly	Pro	Gly	Ser	Pro	Glu	Ser	Ile	Lys	
475															485	
480																
1824																
GCA	TAC	CTT	AAA	CAA	GCT	CCA	CCT	TCC	AAA	GGA	CCT	ACT	GTT	CGG	ACA	1824
Ala	Tyr	Leu	Lys	Gln	Ala	Pro	Pro	Ser	Lys	Gly	Pro	Thr	Val	Arg	Thr	
490															500	
495															505	
1872																
AAA	AAA	GTA	GGG	AAA	AAC	GAA	GCT	GTC	TTA	GAG	TGG	GAC	CRA	CTT	CCT	1872
Lys	Lys	Val	Gly	Lys	Asn	Glu	Ala	Val	Leu	Glu	Trp	Asp	Gln	Leu	Pro	
510															520	
515																
1920																
GTT	GAT	GTT	CAG	AAT	GGA	TTT	ATC	AGA	AAT	TAT	ACT	ATA	TTT	TAT	AGA	1920
Val	Asp	Val	Gln	Asn	Gly	Phe	Ile	Arg	Asn	Tyr	Thr	Ile	Phe	Tyr	Arg	
525															535	
530																
1968																
ACC	ATC	ATT	GGA	AAT	GAA	ACT	GCT	GTG	AAT	GTG	GAT	TCT	TCC	CAC	ACA	1968
Thr	Ile	Ile	Gly	Asn	Glu	Thr	Ala	Val	Asn	Val	Asp	Ser	Ser	His	Thr	
540															550	
545																
2016																
GAA	TAT	ACA	TTG	TCC	TCT	TTG	ACT	AGT	GAC	ACA	TTG	TAC	ATG	GTA	CGA	2016
Glu	Tyr	Thr	Leu	Ser	Ser	Leu	Thr	Ser	Asp	Thr	Leu	Tyr	Met	Val	Arg	
555															565	
560																
2064																
AT																

WO 93/10151

PCT/US92/10272

51

CCT	GTT	TGC	TTA	GCA	TTC	CTA	TTG	ACA	ACT	CTT	CTG	GGA	GTG	CTG	TTC	2160
Pro	Val	Cys	Leu	Ala	Phe	Leu	Leu	Thr	Thr	Leu	Leu	Gly	Val	Leu	Phe	
			605					610					615			
TGC	TTT	AAT	AAG	CGA	GAC	CTA	ATT	AAA	AAA	CAC	ATC	TGG	CCT	AAT	GTT	2208
Cys	Phe	Asn	Lys	Arg	Asp	Leu	Ile	Lys	Lys	His	Ile	Trp	Pro	Asn	Val	
		620					625					630				
CCA	GAT	CCT	TCA	AAG	AGT	CAT	ATT	GCC	CAG	TGG	TCA	CCT	CAC	ACT	CCT	2256
Pro	Asp	Pro	Ser	Lys	Ser	His	Ile	Ala	Gln	Trp	Ser	Pro	His	Thr	Pro	
	635					640					645					
CCA	AGG	CAC	AAT	TTT	AAT	TCA	AAA	GAT	CAA	ATG	TAT	TCA	GAT	GGC	AAT	2304
Pro	Arg	His	Asn	Phe	Asn	Ser	Lys	Asp	Gln	Met	Tyr	Ser	Asp	Gly	Asn	
650					655				660						665	
TTC	ACT	GAT	GTA	AGT	GTT	GTG	GAA	ATA	GAA	GCA	AAT	GAC	AAA	AAG	CCT	2352
Phe	Thr	Asp	Val	Ser	Val	Val	Glu	Ile	Glu	Ala	Asn	Asp	Lys	Lys	Pro	
				670				675						680		
TTT	CCA	GAA	GAT	CTG	AA											2369
Phe	Pro	Glu	Asp	Leu												
			685													

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 708 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Leu	Thr	Leu	Gln	Thr	Trp	Leu	Val	Gln	Ala	Leu	Phe	Ile	Phe	Leu	
-22		-20					-15					-10				
Thr	Thr	Glu	Ser	Thr	Gly	Glu	Leu	Leu	Asp	Pro	Cys	Gly	Tyr	Ile	Ser	
-5						1				5					10	
Pro	Glu	Ser	Pro	Val	Val	Gln	Leu	His	Ser	Asn	Phe	Thr	Ala	Val	Cys	
				15					20					25		
Val	Leu	Lys	Glu	Lys	Cys	Met	Asp	Tyr	Phe	His	Val	Asn	Ala	Asn	Tyr	
		30					35						40			
Ile	Val	Trp	Lys	Thr	Asn	His	Phe	Thr	Ile	Pro	Lys	Glu	Gln	Tyr	Thr	
		45					50					55				
Ile	Ile	Asn	Arg	Thr	Ala	Ser	Ser	Val	Thr	Phe	Thr	Asp	Ile	Ala	Ser	
	60					65					70					
Leu	Asn	Ile	Gln	Leu	Thr	Cys	Asn	Ile	Leu	Thr	Phe	Gly	Gln	Leu	Glu	
	75				80					85					90	
Gln	Asn	Val	Tyr	Gly	Ile	Thr	Ile	Ile	Ser	Gly	Leu	Pro	Pro	Glu	Lys	
				95					100						105	

WO 93/10151

PCT/US92/10272

52

Pro Lys Asn Leu Ser Cys Ile Val Asn Glu Gly Lys^c Lys Met Arg Cys
 110 115 120

Glu Trp Asp Gly Gly Arg Glu Thr His Leu Glu Thr Asn Phe Thr Leu
 125 130 135

Lys Ser Glu Trp Ala Thr His Lys Phe Ala Asp Cys Lys Ala Lys Arg
 140 145 150

Asp Thr Pro Thr Ser Cys Thr Val Asp Tyr Ser Thr Val Tyr Phe Val
 155 160 165 170

Asn Ile Glu Val Trp Val Glu Ala Glu Asn Ala Leu Gly Lys Val Thr
 175 180 185

Ser Asp His Ile Asn Phe Asp Pro Val Tyr Lys Val Lys Pro Asn Pro
 190 195 200

Pro His Asn Leu Ser Val Ile Asn Ser Glu Glu Leu Ser Ser Ile Leu
 205 210 215

Lys Leu Thr Trp Thr Asn Pro Ser Ile Lys Ser Val Ile Ile Leu Lys
 220 225 230

Tyr Asn Ile Gln Tyr Arg Thr Lys Asp Ala Ser Thr Trp Ser Gln Ile
 235 240 245 250

Pro Pro Glu Asp Thr Ala Ser Thr Arg Ser Ser Phe Thr Val Gln Asp
 255 260 265

Leu Lys Pro Phe Thr Glu Tyr Val Phe Arg Ile Arg Cys Met Lys Glu
 270 275 280

Asp Gly Lys Gly Tyr Trp Ser Asp Trp Ser Glu Glu Ala Ser Gly Ile
 285 290 295

Thr Tyr Glu Asp Arg Pro Ser Lys Ala Pro Ser Phe Trp Tyr Lys Ile
 300 305 310

Asp Pro Ser His Thr Gln Gly Tyr Arg Thr Val Gln Leu Val Trp Lys
 315 320 325 330

Thr Leu Pro Pro Phe Gln Ala Asn Gly Lys Ile Leu Asp Tyr Glu Val
 335 340 345

Thr Leu Thr Arg Trp Lys Ser His Leu Gln Asn Tyr Thr Val Asn Ala
 350 355 360

Thr Lys Leu Thr Val Asn Leu Thr Asn Asp Arg Tyr Leu Ala Thr Leu
 365 370 375

Thr Val Arg Asn Leu Val Gly Lys Ser Asp Ala Ala Val Leu Thr Ile
 380 385 390

Pro Ala Cys Asp Phe Gln Ala Thr His Pro Val Met Asp Leu Lys Ala
 395 400 405 410

Phe Pro Lys Asp Asn Met Leu Trp Val Glu Trp Thr Thr Pro Arg Glu
 415 420 425

WO 93/10131

PCT/US92/10272

53

Ser Val Lys Lys Tyr Ile Leu Glu Trp Cys Val Leu Ser Asp Lys Ala
 430 435 440
 Pro Cys Ile Thr Asp Trp Gln Gln Glu Asp Gly Thr Val His Arg Thr
 445 450 455
 Tyr Leu Arg Gly Asn Leu Ala Glu Ser Lys Cys Tyr Leu Ile Thr Val
 460 465 470
 Thr Pro Val Tyr Ala Asp Gly Pro Gly Ser Pro Glu Ser Ile Lys Ala
 475 480 485 490
 Tyr Leu Lys Gln Ala Pro Pro Ser Lys Gly Pro Thr Val Arg Thr Lys
 495 500 505
 Lys Val Gly Lys Asn Glu Ala Val Leu Glu Trp Asp Gln Leu Pro Val
 510 515 520
 Asp Val Gln Asn Gly Phe Ile Arg Asn Tyr Thr Ile Phe Tyr Arg Thr
 525 530 535
 Ile Ile Gly Asn Glu Thr Ala Val Asn Val Asp Ser Ser His Thr Glu
 540 545 550
 Tyr Thr Leu Ser Ser Leu Thr Ser Asp Thr Leu Tyr Met Val Arg Met
 555 560 565 570
 Ala Ala Tyr Thr Asp Glu Gly Gly Lys Asp Gly Pro Glu Phe Thr Phe
 575 580 585
 Thr Thr Pro Lys Phe Ala Gln Gly Glu Ile Glu Ala Ile Val Val Pro
 590 595 600
 Val Cys Leu Ala Phe Leu Leu Thr Thr Leu Leu Gly Val Leu Phe Cys
 605 610 615
 Phe Asn Lys Arg Asp Leu Ile Lys Lys His Ile Trp Pro Asn Val Pro
 620 625 630
 Asp Pro Ser Lys Ser His Ile Ala Gln Trp Ser Pro His Thr Pro Pro
 635 640 645 650
 Arg His Asn Phe Asn Ser Lys Asp Gln Met Tyr Ser Asp Gly Asn Phe
 655 660 665
 Thr Asp Val Ser Val Val Glu Ile Glu Ala Asn Asp Lys Lys Pro Phe
 670 675 680
 Pro Glu Asp Leu
 685

(2) INFORMATION FOR SEQ ID NO:3:

- (1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 705 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(11) MOLECULE TYPE: cDNA to mRNA

PCT/US92/10272

WO 93/10151

54

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:

(B) CLONE: hlgG1Fc

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..699

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GAG CCC AGA TCT TGT GAC AAA ACT CAC ACA TGC CCA CCG TGC CCA GCA	48
Glu Pro Arg Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala	
1 5 10 15	
CCT GAA CTC CTG GGG GGA CCG TCA GTC TTC CTC TTC CCC CCA AAA CCC	96
Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro	
20 25 30	
AAG GAC ACC CTC ATG ATC TCC CGG ACC CCT GAG GTC ACA TGC GTG GTG	144
Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val	
35 40 45	
GTG GAC GTG AGC CAC GAA GAC CCT GAG GTC AAG TTC AAC TGG TAC GTG	192
Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val	
50 55 60	
GAC GGC GTG GAG GTG CAT AAT GCC AAG ACA AAG CCG CGG GAG GAG CAG	240
Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Glu	
65 70 75 80	
EAC AAC AGC ACG TAC CGG GTG GTC AGC GTC CTC ACC GTC CTC CAC CAG	288
Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Glu	
85 90 95	
GAC TGG CTG AAT GGC AAG GAC TAC AAG TGC AAG GTC TCC AAC AAA GCC	336
Asp Trp Leu Asn Gly Lys Asp Tyr Lys Cys Lys Val Ser Asn Lys Ala	
100 105 110	
CTC CCA GCC CCC ATG CAG AAA ACC ATC TCC AAA GCC AAA GGG CAG CCC	384
Leu Pro Ala Pro Met Glu Lys Thr Ile Ser Lys Ala Lys Gly Glu Pro	
115 120 125	
CGA GAA CCA CAG GTG TAC ACC CTG CCC CCA TCC CGG GAT GAG CTG ACC	432
Arg Glu Pro Glu Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr	
130 135 140	
AAG AAC CAG GTC AGC CTG ACC TGC CTG GTC AAA GGC TTC TAT CCC AGG	480
Lys Asn Glu Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Arg	
145 150 155 160	
CAC ATC GCC GTG GAG TGG GAG AGC AAT GCG CAG CCG GAG AAC AAC TAC	528
His Ile Ala Val Glu Trp Glu Ser Asn Gly Glu Pro Glu Asn Asn Tyr	
165 170 175	

WO 93/10151

PCT/US92/10272

55

AAG ACC ACG OCT CCC GTG CTG GAC TCC GAC GGC TCC TTC TTC CTC TAC	576
Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr	
180 185 190	
AGC AAG CTC ACC GTG GAC AAG AGC AGG TGG CAG CAG GGG AAC GTC TTC	624
Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe	
195 200 205	
TCA TGC TCC GTG ATG CAT GAG GCT CTG CAC AAC CAC TAC ACG CAG AAG	672
Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys	
210 215 220	
AGC CTC TCC CTG TCT CCG GGT AAA TGA ACT AGT	705
Ser Leu Ser Leu Ser Pro Gly Lys	
225 230	

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 232 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Glu	Pro	Arg	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala
1				5					10					15	
Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro
		20						25					30		
Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val
		35					40					45			
Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val
		50				55					60				
Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln
65					70					75					80
Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln
				85					90					95	
Asp	Trp	Leu	Asn	Gly	Lys	Asp	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala
		100						105					110		
Leu	Pro	Ala	Pro	Met	Gln	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro
		115					120					125			
Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr
		130				135					140				
Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Arg
145					150					155					160
His	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr
				165					170					175	

WO 93/10151

PCT/US92/10272

[illegible]

(2) INFORMATION FOR SEQ ID NO:5:

(1) SEQUENCE CHARACTERISTICS:

- SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 3182 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(11) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: N-terminal

(vi) ORIGINAL SOURCE:

- (F) TISSUE TYPE: human placenta

(V11) IMMEDIATE SOURCE:

- (E) CLONE: PHLIFR-65

(ix) **FEATURE:**

- ```
FEATURE:
(A) NAME/KEY: mat_peptide
(B) LOCATION: 311..3182
```

(ix) **FEATURE:**

- (A) NAME/KEY: CDS  
(B) LOCATION: 179..3182

(1x) FEATURE:

- FEATURE:  
(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 179..310

(ix) **FEATURE:**

- FEATURE:
- (A) NAME/KEY: mat\_peptide
- (B) LOCATION: 311..3182

(~~2~~) SEQUENCE DESCRIPTION: SEQ ID NO:5:

| SEQUENCE POSITION | SEQUENCE   | POSITION    |            |            |            |     |
|-------------------|------------|-------------|------------|------------|------------|-----|
| AGATCTTGGA        | ACGAGACGAC | CTGCTCTCTC  | TCCAGAACG  | TGTCTCTGCT | GCAAGGCACC | 60  |
| GGGCCCCCTTC       | GCTCTGCAGA | ACTGCACTTG  | CAAGACCATT | ATCAACTCCT | AATCCAGCT  | 120 |
| CAGAAAGGGA        | GCCTCTGCGA | CTCATTTCATC | GCCCTCCAGG | ACTGACTGCA | TTGCACAG   | 178 |

WO 93/10151

PCT/US92/10272

57

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| ATG | ATG | GAT | ATT | TAC | GTA | TGT | TTG | AAA | CGA | CCA | TCC | TGG | ATG | GTG | GAC | 226 |
| Met | Met | Asp | Ile | Tyr | Val | Cys | Leu | Lys | Arg | Pro | Ser | Trp | Met | Val | Asp |     |
| -44 |     |     |     | -40 |     |     |     |     | -35 |     |     |     |     | -30 |     |     |
| AAT | AAA | AGA | ATG | AGG | ACT | GCT | TCA | AAT | TTC | CAG | TGG | CTG | TTA | TCA | ACA | 274 |
| Asn | Lys | Arg | Met | Arg | Thr | Ala | Ser | Asn | Phe | Gln | Trp | Leu | Leu | Ser | Thr |     |
|     |     |     | -25 |     |     |     |     | -20 |     |     |     |     | -15 |     |     |     |
| TTT | ATT | CTT | CTA | TAT | CTA | ATG | AAT | CAA | GTA | AAT | AGC | CAG | AAA | AAG | GGG | 322 |
| Phe | Ile | Leu | Leu | Tyr | Leu | Met | Asn | Gln | Val | Asn | Ser | Gln | Lys | Lys | Gly |     |
|     |     | -10 |     |     |     |     | -5  |     |     |     |     | 1   |     |     |     |     |
| GCT | CCT | CAT | GAT | TTG | AAG | TGT | GTA | ACT | AAC | AAT | TTG | CAA | GTG | TGG | AAC | 370 |
| Ala | Pro | His | Asp | Leu | Lys | Cys | Val | Thr | Asn | Asn | Leu | Gln | Val | Trp | Asn |     |
| 6   |     |     |     | 10  |     |     |     |     | 15  |     |     |     |     | 20  |     |     |
| TGT | TCT | TGG | AAA | GCA | CCC | TCT | GGA | ACA | GGC | CGT | GGT | ACT | GAT | TAT | GAA | 418 |
| Cys | Ser | Trp | Lys | Ala | Pro | Ser | Gly | Thr | Gly | Arg | Gly | Thr | Asp | Tyr | Glu |     |
|     |     |     | 25  |     |     |     |     |     | 30  |     |     |     |     | 35  |     |     |
| GTT | TGC | ATT | GAA | AAC | AGG | TCC | CGT | TCT | TGT | TAT | CAG | TTG | GAG | AAA | ACC | 466 |
| Val | Cys | Ile | Glu | Asn | Arg | Ser | Arg | Ser | Cys | Tyr | Gln | Leu | Glu | Lys | Thr |     |
|     |     |     | 40  |     |     |     |     | 45  |     |     |     |     | 50  |     |     |     |
| AGT | ATT | AAA | ATT | CCA | GCT | CTT | TCA | CAT | GGT | GAT | TAT | GAA | ATA | ACA | ATA | 514 |
| Ser | Ile | Lys | Ile | Pro | Ala | Leu | Ser | His | Gly | Asp | Tyr | Glu | Ile | Thr | Ile |     |
|     |     | 35  |     |     |     |     | 60  |     |     |     |     | 65  |     |     |     |     |
| AAT | TCT | CTA | CAT | GAT | TTT | GGA | AGT | TCT | ACA | AGT | AAA | TTC | ACA | CTA | AAT | 562 |
| Asn | Ser | Leu | His | Asp | Phe | Gly | Ser | Ser | Thr | Ser | Lys | Phe | Thr | Leu | Asn |     |
|     | 70  |     |     |     |     | 75  |     |     |     |     | 80  |     |     |     |     |     |
| GAA | CAA | AAC | GTT | TCC | TTA | ATT | CCA | GAT | ACT | CCA | GAG | ATC | TTG | AAT | TTG | 610 |
| Glu | Gln | Asn | Val | Ser | Leu | Ile | Pro | Asp | Thr | Pro | Glu | Ile | Leu | Asn | Leu |     |
| 85  |     |     |     |     | 90  |     |     |     |     | 95  |     |     |     |     | 100 |     |
| TCT | GCT | GAT | TTC | TCA | ACC | TCT | ACA | TTA | TAC | CTA | AAG | TGG | AAC | GAC | AGG | 638 |
| Ser | Ala | Asp | Phe | Ser | Thr | Ser | Thr | Leu | Tyr | Leu | Lys | Trp | Asn | Asp | Arg |     |
|     |     |     | 105 |     |     |     |     |     | 110 |     |     |     |     | 115 |     |     |
| GGT | TCA | GTT | TTT | CCA | CAC | CGC | TCA | AAT | GTT | ATC | TGG | GAA | ATT | AAA | GTT | 706 |
| Gly | Ser | Val | Phe | Pro | His | Arg | Ser | Asn | Val | Ile | Trp | Glu | Ile | Lys | Val |     |
|     |     |     | 120 |     |     |     |     | 125 |     |     |     |     | 130 |     |     |     |
| CTA | CGT | AAA | GAG | AGT | ATG | GAG | CTC | GTA | AAA | TTA | GTG | ACC | CAC | AAC | ACA | 754 |
| Leu | Arg | Lys | Glu | Ser | Met | Glu | Leu | Val | Lys | Leu | Val | Thr | His | Asn | Thr |     |
|     |     | 135 |     |     |     |     | 140 |     |     |     |     | 145 |     |     |     |     |
| ACT | CTG | AAT | GGC | AAA | GAT | ACA | CTT | CAT | CAC | TGG | AGT | TGG | GCC | TCA | GAT | 802 |
| Thr | Leu | Asn | Gly | Lys | Asp | Thr | Leu | His | His | Trp | Ser | Trp | Ala | Ser | Asp |     |
|     |     | 150 |     |     |     | 155 |     |     |     |     | 160 |     |     |     |     |     |
| ATG | CCC | TTG | GAA | TGT | GCC | ATT | CAT | TTT | GTG | GAA | ATT | AGA | TGC | TAC | ATT | 850 |
| Met | Pro | Leu | Glu | Cys | Ala | Ile | His | Phe | Val | Glu | Ile | Arg | Cys | Tyr | Ile |     |
| 165 |     |     |     | 170 |     |     |     |     | 175 |     |     |     |     | 180 |     |     |
| GAC | AAT | CTT | CAT | TTT | TCT | GGT | CTC | GAA | GAG | TGG | AGT | GAC | TGG | AGC | CCT | 898 |
| Asp | Asn | Leu | His | Phe | Ser | Gly | Leu | Glu | Glu | Trp | Ser | Asp | Trp | Ser | Pro |     |
|     |     |     | 185 |     |     |     |     | 190 |     |     |     |     |     | 195 |     |     |

**PCT/US92/10272**

| 58  |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     | 946  |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|
| GTG | AAG | AAC | ATT | TCT | TGG | ATA | CCT | GAT | TCT | CAG | ACT | AAG | GTT | TTT | CCT | 946  |
| Val | Lys | Asn | Ile | Ser | Trp | Ile | Pro | Asp | Ser | Gln | Thr | Lys | Val | Phe | Pro |      |
| 200 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     | 210  |
| CAA | GAT | AAA | GTG | ATA | CTT | GTA | GGC | TCA | GAC | ATA | ACA | TTT | TGT | TGT | GTG | 994  |
| Gln | Asp | Lys | Val | Ile | Leu | Val | Gly | Ser | Asp | Ile | Thr | Phe | Cys | Cys | Val |      |
| 215 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     | 225  |
| AGT | CAA | GAA | AAA | GTG | TTA | TCA | GCA | CTG | ATT | GGC | CAT | ACA | AAC | TGC | CCC | 1042 |
| Ser | Gln | Glu | Lys | Val | Leu | Ser | Ala | Leu | Ile | Gly | His | Thr | Asn | Cys | Pro |      |
| 230 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     | 235  |
| TTG | ATC | CAT | CTT | GAT | GGG | GAA | AAT | GTT | GCA | ATC | AAG | ATT | CGT | AAT | ATT | 1090 |
| Leu | Ile | His | Leu | Asp | Gly | Gln | Asn | Val | Ala | Ile | Lys | Ile | Arg | Asn | Ile |      |
| 245 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     | 250  |
| TCT | GTT | TCT | GCA | AGT | AGT | GGA | ACA | AAT | GTA | GTT | TTT | ACA | ACC | GAA | GAT | 1138 |
| Ser | Val | Ser | Ala | Ser | Ser | Gly | Thr | Asn | Val | Val | Phe | Thr | Thr | Glu | Asp |      |
| 265 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     | 270  |
| AAC | ATA | TTT | GGA | ACC | GTT | ATT | TTT | GCT | GGA | TAT | CCA | CCA | GAT | ACT | CCT | 1186 |
| Asn | Ile | Phe | Gly | Thr | Val | Ile | Phe | Ala | Gly | Tyr | Pro | Pro | Asp | Thr | Pro |      |
| 280 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     | 285  |
| CAA | CAA | CTG | AAT | TGT | GAG | ACA | CAT | GAT | TTA | AAA | GAA | ATT | ATA | TGT | AGT | 1234 |
| Gln | Gln | Leu | Asn | Cys | Gln | Thr | His | Asp | Leu | Lys | Glu | Ile | Ile | Cys | Ser |      |
| 295 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     | 300  |
| TGG | AAT | CCA | GGA | AGG | GTG | ACA | CGG | TTG | GTG | GGC | CCA | CGT | GCT | ACA | AGC | 1282 |
| Trp | Asn | Pro | Gly | Arg | Val | Thr | Ala | Leu | Val | Gly | Pro | Arg | Ala | Thr | Ser |      |
| 310 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     | 315  |
| TAC | ACT | TTA | GTT | GAA | AGT | TTT | TCA | GGA | AAA | TAT | GTT | AGA | CTT | AAA | AGA | 1330 |
| Tyr | Thr | Leu | Val | Glu | Ser | Phe | Ser | Gly | Lys | Tyr | Val | Arg | Leu | Lys | Arg |      |
| 325 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     | 330  |
| GCT | GAA | GCA | CCT | ACA | AAC | GAA | AGC | TAT | CAA | TTA | TTA | TTT | CAA | ATG | CTT | 1378 |
| Ala | Glu | Ala | Pro | Thr | Asn | Glu | Ser | Tyr | Gln | Leu | Leu | Phe | Gln | Met | Leu |      |
| 345 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     | 350  |
| CCA | AAT | CAA | GAA | ATA | TAT | AAT | TTT | ACT | TTG | AAT | GCT | CAC | AAT | CCG | CTG | 1426 |
| Pro | Asn | Gln | Glu | Ile | Tyr | Asn | Phe | Thr | Leu | Asn | Ala | His | Asn | Pro | Leu |      |
| 360 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     | 365  |
| GGT | CGA | TCA | CAA | TCA | ACA | ATT | TTA | GTT | AAT | ATA | ACT | GAA | AAA | GTT | TAT | 1474 |
| Gly | Arg | Ser | Gln | Ser | Thr | Ile | Leu | Val | Asn | Ile | Thr | Glu | Lys | Val | Tyr |      |
| 375 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     | 380  |
| CCC | CAT | ACT | CCT | ACT | TCA | TTC | AAA | GTG | AAG | GAT | ATT | AAT | TCA | ACA | GCT | 1522 |
| Pro | His | Thr | Pro | Thr | Ser | Phe | Lys | Val | Lys | Asp | Ile | Asn | Ser | Thr | Ala |      |
| 390 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     | 395  |
| GTT | AAA | CTT | TCT | TGG | CAT | TTA | CCA | GGC | AAC | TTT | GCA | AAG | ATT | AAT | TTT | 1570 |
| Val | Lys | Leu | Ser | Trp | His | Leu | Pro | Gly | Asn | Phe | Ala | Lys | Ile | Asn | Phe |      |
| 405 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     | 410  |
| TTA | TGT | GAA | ATT | GAA | ATT | AAG | AAA | TCT | AAT | TCA | GTA | CAA | GAG | CAG | CGG | 1618 |
| Leu | Cys | Glu | Ile | Gln | Ile | Lys | Lys | Ser | Asn | Ser | Val | Gln | Gln | Gln | Arg |      |
| 425 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     | 430  |

WO 93/10151

PCT/US92/10272

59

|                                                                                                                                                       |      |
|-------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| AAT GTC ACA ATC AAA GGA GTA GAA AAT TCA AGT TAT CTT GTT GCT CTG<br>Asn Val Thr Ile Lys Gly Val Glu Asn Ser Ser Tyr Leu Val Ala Leu<br>440 445 450     | 1666 |
| GAC AAG TTA AAT CCA TAC ACT CTA TAT ACT TTT CGG ATT CGT TGT TCT<br>Asp Lys Leu Asn Pro Tyr Thr Leu Tyr Thr Phe Arg Ile Arg Cys Ser<br>455 460 465     | 1714 |
| ACT GAA ACT TTC TGG AAA TGG AGC AAA TGG AGC AAT AAA AAA CAA CAT<br>Thr Glu Thr Phe Trp Lys Trp Ser Lys Trp Ser Asn Lys Lys Gln His<br>470 475 480     | 1762 |
| TTA ACA ACA GAA GCC AGT CCT TCA AAG GGG CCT GAT ACT TGG AGA GAG<br>Leu Thr Thr Glu Ala Ser Pro Ser Lys Gly Pro Asp Thr Trp Arg Glu<br>485 490 495 500 | 1810 |
| TGG AGT TCT GAT GGA AAA AAT TTA ATA ATC TAT TGG AAG CCT TTA CCC<br>Trp Ser Ser Asp Gly Lys Asn Leu Ile Ile Tyr Trp Lys Pro Leu Pro<br>505 510 515     | 1858 |
| ATT AAT GAA CCT AAT GGA AAA ATA CTT TCC TAC AAT GTA TCG TGT TCA<br>Ile Asn Gln Ala Asn Gly Lys Ile Leu Ser Tyr Asn Val Ser Cys Ser<br>520 525 530     | 1906 |
| TCA GAT GAG GAA ACA CAG TCC CTT TCT GAA ATC CCT GAT CCT CAG CAC<br>Ser Asp Glu Glu Thr Gln Ser Leu Ser Glu Ile Pro Asp Pro Gln His<br>535 540 545     | 1954 |
| AAA GCA GAG ATA CGA CTT GAT AAG AAT GAC TAC ATC ATC AGC GTA GTG<br>Lys Ala Glu Ile Arg Leu Asp Lys Asn Asp Tyr Ile Ile Ser Val Val<br>550 555 560     | 2002 |
| GCT AAA AAT TCT GTG GGC TCA TCA CCA CCT TCC AAA ATA GCG AGT ATG<br>Ala Lys Asn Ser Val Gly Ser Ser Pro Pro Ser Lys Ile Ala Ser Met<br>565 570 575 580 | 2050 |
| GAA ATT CCA AAT GAT GAT CTC AAA ATA GAA CAA GTT GTT GGG ATG GGA<br>Glu Ile Pro Asn Asp Asp Leu Lys Ile Glu Gln Val Val Gly Met Gly<br>585 590 595     | 2098 |
| AAG GGG ATT CTC CTC ACC TGG CAT TAC GAC CCC AAC ATG ACT TGC GAC<br>Lys Gly Ile Leu Leu Thr Trp His Tyr Asp Pro Asn Met Thr Cys Asp<br>600 605 610     | 2146 |
| TAC GTC ATT AAG TGG TGT AAC TCG TCT CGG TCG GAA CCA TGC CTT ATG<br>Tyr Val Ile Lys Trp Cys Asn Ser Ser Arg Ser Glu Pro Cys Leu Met<br>615 620 625     | 2194 |
| GAC TGG AGA AAA GTT CCC TCA AAC AGC ACT GAA ACT GTA ATA GAA TCT<br>Asp Trp Arg Lys Val Pro Ser Asn Ser Thr Glu Thr Val Ile Glu Ser<br>630 635 640     | 2242 |
| GAT GAG TTT CGA CCA GGT ATA AGA TAT AAT TTT TTC CTG TAT GGA TGC<br>Asp Glu Phe Arg Pro Gly Ile Arg Tyr Asn Phe Phe Leu Tyr Gly Cys<br>645 650 655 660 | 2290 |
| AGA AAT CAA GGA TAT CAA TTA TTA CGC TCC ATG ATT GGA TAT ATA GAA<br>Arg Asn Gln Gly Tyr Gln Leu Leu Arg Ser Met Ile Gly Tyr Ile Glu<br>665 670 675     | 2338 |

PCT/US92/10272

WO 93/10151

60

|                                                                 |      |
|-----------------------------------------------------------------|------|
| GAA TTG GCT CCC ATT GTT GCA CCA AAT TTT ACT GTT GAG GAT ACT TCT | 2386 |
| Glu Leu Ala Pro Ile Val Ala Pro Asn Phe Thr Val Glu Asp Thr Ser |      |
| 680 685 690                                                     |      |
| GCA GAT TCG ATA TTA GTA ABA TGG GAA GAC ATT CCT GTG GAA GAA CTT | 2434 |
| Ala Asp Ser Ile Leu Val Lys Trp Glu Asp Ile Pro Val Glu Glu Leu |      |
| 695 700 705                                                     |      |
| AGA GGC TTT TTA AGA GGA TAT TTG TTT TAC TTT GGA AAA CGA GAA AGA | 2482 |
| Arg Gly Phe Leu Arg Gly Tyr Leu Phe Tyr Phe Gly Lys Gly Glu Arg |      |
| 710 715 720                                                     |      |
| GAC ACA TCT AAG ATG AGG GTT TTA GAA TCA GGT CGT TCT GAC ATA AAA | 2530 |
| Asp Thr Ser Lys Met Arg Val Leu Glu Ser Gly Arg Ser Asp Ile Lys |      |
| 725 730 735 740                                                 |      |
| GTT AAG AAT ATT ACT GAC ATA TCC CAG AAG ACA CTG AGA ATT GCT GAT | 2578 |
| Val Lys Asn Ile Thr Asp Ile Ser Gln Lys Thr Leu Arg Ile Ala Asp |      |
| 745 750 755                                                     |      |
| CTT CAA GGT AAA ACA AGT TAC CAC CTG GTC TTG CGA GCC TAT ACA GAT | 2626 |
| Leu Gln Gly Lys Thr Ser Tyr His Leu Val Leu Arg Ala Tyr Thr Asp |      |
| 760 765 770                                                     |      |
| GGT GGA GTG GGC CCG GAG AAG AGT ATG TAT GTG GTG ACA AAG GAA AAT | 2674 |
| Gly Gly Val Gly Pro Glu Lys Ser Met Tyr Val Val Thr Lys Glu Asn |      |
| 775 780 785                                                     |      |
| TCT GTG GGA TTA ATT ATT GCC ATT CTC ATC CCA GTG GCA GTG GCT GTC | 2722 |
| Ser Val Gly Leu Ile Ile Ala Ile Leu Ile Pro Val Ala Val Ala Val |      |
| 790 795 800                                                     |      |
| ATT GTT GGA GTG GTG ACA AGT ATC CTT TGC TAT CGG AAA CGB GAA TGG | 2770 |
| Ile Val Gly Val Val Thr Ser Ile Leu Cys Tyr Arg Lys Arg Glu Trp |      |
| 805 810 815 820                                                 |      |
| ATT AAA GAA ACC TTC TAC CCT GAT ATT CCA AAT CCA GAA AAC TGT AAA | 2818 |
| Ile Lys Glu Thr Phe Tyr Pro Asp Ile Pro Asn Pro Glu Asn Cys Lys |      |
| 825 830 835                                                     |      |
| GCA TTA CAG TTT CAA AAG AGT GTC TGT GAG GGA AGC AGT GCT CTT ABA | 2866 |
| Ala Leu Gln Phe Gln Lys Ser Val Cys Glu Gly Ser Ser Ala Leu Lys |      |
| 840 845 850                                                     |      |
| ACA TTG GAA ATG AAT CCT TGT ACC CCA AAT AAT GTT GAG GTT CTG GAA | 2914 |
| Thr Leu Glu Met Asn Pro Cys Thr Pro Asn Asn Val Glu Val Leu Glu |      |
| 855 860 865                                                     |      |
| ACT CGA TCA GCA TTT CCT AAA ATA GAA GAT ACA GAA ATA ATT TCC CCA | 2962 |
| Thr Arg Ser Ala Phe Pro Lys Ile Glu Asp Thr Glu Ile Ile Ser Pro |      |
| 870 875 880                                                     |      |
| GTA GCT GAG CGT CCT GAA GAT CGC TCT GAT GCA GAG CCT GAA AAC CAT | 3010 |
| Val Ala Glu Arg Pro Glu Asp Arg Ser Asp Ala Glu Pro Glu Asn His |      |
| 885 890 895 900                                                 |      |
| GTG GTT GTG TCC TAT TGT CCA CCC ATC ATT GAG GAA GAA ATA CCA AAC | 3058 |
| Val Val Val Ser Tyr Cys Pro Pro Ile Ile Glu Glu Glu Ile Pro Asn |      |
| 905 910 915                                                     |      |

WO 93/10151

PCT/US92/10272

61

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |      |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|
| CCA | GCC | GCA | GAT | GAA | GCT | GGA | GGG | ACT | GCA | CAG | GTT | ATT | TAC | ATT | GAT | 3106 |
| Pro | Ala | Ala | Asp | Glu | Ala | Gly | Gly | Thr | Ala | Gln | Val | Ile | Tyr | Ile | Asp |      |
|     |     |     | 920 |     |     |     |     | 925 |     |     |     |     | 930 |     |     |      |
|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |      |
| GTT | CAG | TCG | ATG | TAT | CAG | CCT | CAA | GCA | AAA | CCA | GAA | GAA | AAA | AAA | AAA | 3154 |
| Val | Gln | Ser | Met | Tyr | Gln | Pro | Gln | Ala | Lys | Pro | Glu | Glu | Lys | Lys | Lys |      |
|     |     | 935 |     |     |     |     | 940 |     |     |     |     | 945 |     |     |     |      |
|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |      |
| AAA | AGC | AGG | TCG | TCT | CGT | TCC | AAG | ATC | T   |     |     |     |     |     |     | 3182 |
| Lys | Ser | Arg | Ser | Ser | Arg | Ser | Lys | Ile |     |     |     |     |     |     |     |      |
|     |     | 950 |     |     |     |     | 955 |     |     |     |     |     |     |     |     |      |

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1001 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|
| Met | Met | Asp | Ile | Tyr | Val | Cys | Leu | Lys | Arg | Pro | Ser | Trp | Met | Val | Asp |  |
| -44 |     |     |     | -40 |     |     |     |     | -35 |     |     |     |     | -30 |     |  |
|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  |
| Asn | Lys | Arg | Met | Arg | Thr | Ala | Ser | Asn | Phe | Gln | Trp | Leu | Leu | Ser | Thr |  |
|     |     | -25 |     |     |     |     |     | -20 |     |     |     |     | -15 |     |     |  |
|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  |
| Phe | Ile | Leu | Leu | Tyr | Leu | Met | Asn | Gln | Val | Asn | Ser | Gln | Lys | Lys | Gly |  |
|     |     | -10 |     |     |     | -5  |     |     |     |     |     | 1   |     |     |     |  |
|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  |
| Ala | Pro | His | Asp | Leu | Lys | Cys | Val | Thr | Asn | Asn | Leu | Gln | Val | Trp | Asn |  |
| 5   |     |     |     | 10  |     |     |     |     |     | 15  |     |     |     |     | 20  |  |
|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  |
| Cys | Ser | Trp | Lys | Ala | Pro | Ser | Gly | Thr | Gly | Arg | Gly | Thr | Asp | Tyr | Glu |  |
|     |     |     | 25  |     |     |     |     |     | 30  |     |     |     |     | 35  |     |  |
|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  |
| Val | Cys | Ile | Glu | Asn | Arg | Ser | Arg | Ser | Cys | Tyr | Gln | Leu | Glu | Lys | Thr |  |
|     |     |     | 40  |     |     |     |     | 45  |     |     |     |     | 50  |     |     |  |
|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  |
| Ser | Ile | Lys | Ile | Pro | Ala | Leu | Ser | His | Gly | Asp | Tyr | Glu | Ile | Thr | Ile |  |
|     |     | 55  |     |     |     | 60  |     |     |     |     |     | 65  |     |     |     |  |
|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  |
| Asn | Ser | Leu | His | Asp | Phe | Gly | Ser | Ser | Thr | Ser | Lys | Phe | Thr | Leu | Asn |  |
|     |     | 70  |     |     |     | 75  |     |     |     |     | 80  |     |     |     |     |  |
|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  |
| Glu | Gln | Asn | Val | Ser | Leu | Ile | Pro | Asp | Thr | Pro | Glu | Ile | Leu | Asn | Leu |  |
|     | 85  |     |     |     | 90  |     |     |     |     | 95  |     |     |     |     | 100 |  |
|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  |
| Ser | Ala | Asp | Phe | Ser | Thr | Ser | Thr | Leu | Tyr | Leu | Lys | Trp | Asn | Asp | Arg |  |
|     |     |     | 105 |     |     |     |     |     | 110 |     |     |     |     | 115 |     |  |
|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  |
| Gly | Ser | Val | Phe | Pro | His | Arg | Ser | Asn | Val | Ile | Trp | Glu | Ile | Lys | Val |  |
|     |     |     | 120 |     |     |     |     | 125 |     |     |     |     | 130 |     |     |  |
|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  |
| Leu | Arg | Lys | Glu | Ser | Met | Glu | Leu | Val | Lys | Leu | Val | Thr | His | Asn | Thr |  |
|     |     | 135 |     |     |     |     | 140 |     |     |     |     | 145 |     |     |     |  |

PCT/US92/10272

WO 93/10151

62

Thr Leu Asn Gly Lys Asp Thr Leu His His Trp Ser Trp Ala Ser Asp  
 150 155 160  
 Met Pro Leu Glu Cys Ala Ile His Phe Val Glu Ile Arg Cys Tyr Ile  
 165 170 175 180  
 Asp Asn Leu His Phe Ser Gly Leu Glu Glu Trp Ser Asp Trp Ser Pro  
 185 190 195  
 Val Lys Asn Ile Ser Trp Ile Pro Asp Ser Gln Thr Lys Val Phe Pro  
 200 205 210  
 Gln Asp Lys Val Ile Leu Val Gly Ser Asp Ile Thr Phe Cys Cys Val  
 215 220 225  
 Ser Gln Gln Lys Val Leu Ser Ala Leu Ile Gly His Thr Asn Cys Pro  
 230 235 240  
 Leu Ile His Leu Asp Gly Glu Asn Val Ala Ile Lys Ile Arg Asn Ile  
 245 250 255 260  
 Ser Val Ser Ala Ser Ser Gly Thr Asn Val Val Phe Thr Thr Glu Asp  
 265 270 275  
 Asn Ile Phe Gly Thr Val Ile Phe Ala Gly Tyr Pro Pro Asp Thr Pro  
 280 285 290  
 Gln Gln Leu Asn Cys Glu Thr His Asp Leu Lys Glu Ile Ile Cys Ser  
 295 300 305  
 Trp Asn Pro Gly Arg Val Thr Ala Leu Val Gly Pro Arg Ala Thr Ser  
 310 315 320  
 Tyr Thr Leu Val Gln Ser Phe Ser Gly Lys Tyr Val Arg Leu Lys Arg  
 325 330 335 340  
 Ala Glu Ala Pro Thr Asn Glu Ser Tyr Gln Leu Leu Phe Gln Met Leu  
 345 350 355  
 Pro Asn Gln Glu Ile Tyr Asn Phe Thr Leu Asn Ala His Asn Pro Leu  
 360 365 370  
 Gly Arg Ser Gln Ser Thr Ile Leu Val Asn Ile Thr Glu Lys Val Tyr  
 375 380 385  
 Pro His Thr Pro Thr Ser Phe Lys Val Lys Asp Ile Asn Ser Thr Ala  
 390 395 400  
 Val Lys Leu Ser Trp His Leu Pro Gly Asn Phe Ala Lys Ile Asn Phe  
 405 410 415 420  
 Leu Cys Glu Ile Glu Ile Lys Lys Ser Asn Ser Val Gln Glu Gln Arg  
 425 430 435  
 Asn Val Thr Ile Lys Gly Val Glu Asn Ser Ser Tyr Leu Val Ala Leu  
 440 445 450  
 Asp Lys Leu Asn Pro Tyr Thr Leu Tyr Thr Phe Arg Ile Arg Cys Ser  
 455 460 465

WO 93/10151

PCT/US92/10272

63

Thr Glu Thr Phe Trp Lys Trp Ser Lys Trp Ser Asn Lys Lys Gln His  
 470 475 480  
 Leu Thr Thr Glu Ala Ser Pro Ser Lys Gly Pro Asp Thr Trp Arg Glu  
 485 490 495 500  
 Trp Ser Ser Asp Gly Lys Asn Leu Ile Ile Tyr Trp Lys Pro Leu Pro  
 505 510 515  
 Ile Asn Glu Ala Asn Gly Lys Ile Leu Ser Tyr Asn Val Ser Cys Ser  
 520 525 530  
 Ser Asp Glu Glu Thr Gln Ser Leu Ser Glu Ile Pro Asp Pro Gln His  
 535 540 545  
 Lys Ala Glu Ile Arg Leu Asp Lys Asn Asp Tyr Ile Ile Ser Val Val  
 550 555 560  
 Ala Lys Asn Ser Val Gly Ser Ser Pro Pro Ser Lys Ile Ala Ser Met  
 565 570 575 580  
 Glu Ile Pro Asn Asp Asp Leu Lys Ile Glu Gln Val Val Gly Met Gly  
 585 590 595  
 Lys Gly Ile Leu Leu Thr Trp His Tyr Asp Pro Asn Met Thr Cys Asp  
 600 605 610  
 Tyr Val Ile Lys Trp Cys Asn Ser Ser Arg Ser Glu Pro Cys Leu Met  
 615 620 625  
 Asp Trp Arg Lys Val Pro Ser Asn Ser Thr Glu Thr Val Ile Glu Ser  
 630 635 640  
 Asp Glu Phe Arg Pro Gly Ile Arg Tyr Asn Phe Phe Leu Tyr Gly Cys  
 645 650 655 660  
 Arg Asn Gln Gly Tyr Gln Leu Leu Arg Ser Met Ile Gly Tyr Ile Glu  
 665 670 675  
 Glu Leu Ala Pro Ile Val Ala Pro Asn Phe Thr Val Glu Asp Thr Ser  
 680 685 690  
 Ala Asp Ser Ile Leu Val Lys Trp Glu Asp Ile Pro Val Glu Glu Leu  
 695 700 705  
 Arg Gly Phe Leu Arg Gly Tyr Leu Phe Tyr Phe Gly Lys Gly Glu Arg  
 710 715 720  
 Asp Thr Ser Lys Met Arg Val Leu Glu Ser Gly Arg Ser Asp Ile Lys  
 725 730 735 740  
 Val Lys Asn Ile Thr Asp Ile Ser Gln Lys Thr Leu Arg Ile Ala Asp  
 745 750 755  
 Leu Gln Gly Lys Thr Ser Tyr His Leu Val Leu Arg Ala Tyr Thr Asp  
 760 765 770  
 Gly Gly Val Gly Pro Glu Lys Ser Met Tyr Val Val Thr Lys Glu Asn  
 775 780 785

WO 93/10151

PCI/US92/10272

64

Ser Val Gly Leu Ile Ile Ala Ile Leu Ile Pro Val Ala Val Ala Val  
790 795 800

Ile Val Gly Val Val Thr Ser Ile Leu Cys Tyr Arg Lys Arg Gln Trp  
805 810 815 820

Ile Lys Glu Thr Phe Tyr Pro Asp Ile Pro Asn Pro Glu Asn Cys Lys  
825 830 835

Ala Leu Gln Phe Gln Lys Ser Val Cys Glu Gly Ser Ser Ala Leu Lys  
840 845 850

Thr Leu Glu Met Asn Pro Cys Thr Pro Asn Asn Val Glu Val Leu Glu  
855 860 865

Thr Arg Ser Ala Phe Pro Lys Ile Glu Asp Thr Glu Ile Ile Ser Pro  
870 875 880

Val Ala Glu Arg Pro Glu Asp Arg Ser Asp Ala Glu Pro Glu Asn His  
885 890 895 900

Val Val Val Ser Tyr Cys Pro Pro Ile Ile Glu Glu Glu Ile Pro Asn  
905 910 915

Pro Ala Ala Asp Glu Ala Gly Gly Thr Ala Gln Val Ile Tyr Ile Asp  
920 925 930

Val Gln Ser Met Tyr Gln Pro Gln Ala Lys Pro Gln Glu Lys Lys Lys  
935 940 945

Lys Ser Arg Ser Ser Arg Ser Lys Ile  
950 955

## (2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 100 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GATCCGGTGG AGGTGGTTCT GGTGGAGGTG GTTCAGGTGG TGGAGGATCA GGAGGTGGTG 60

GATCAGGTGG AGGAGGTCTT GGAGGTGGAG GTTCCGGAAT 100

## (2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 36 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

WO 93/10151

PCI/US92/10272

65

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GATATGTCGA CGATGATGCA TATTTACGTA TGTTTG

(2) INFORMATION FOR SEQ ID NO:9:

- (1) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 48 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GCATGGATCC ACCTCCTCCA GAATTTTCCT TTGTCACCAC ATACATAC

(2) INFORMATION FOR SEQ ID NO:10:

- (1) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 49 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CGCGTCCGGA GGAGGTGGAT CTGAACTTCT AGATCCATGT GGTATATC

(2) INFORMATION FOR SEQ ID NO:11:

- (1) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 37 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GCATGCGGCC GCCTATTCAA TTTCTCCTTG ACCAAAC

(2) INFORMATION FOR SEQ ID NO:12:

- (1) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 35 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

WO 93/10151

PCT/US92/10272

66

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GATATGTCGA CAAGATGTTG ACGTTCAGA CTTGG

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 41 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GCATGGATCC ACCCTCTCCT TCAATTCTC CTGAGCAAA C

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 44 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CGCGTCCGGA GGAGGTGCTA GCCAGAAAA GGGGGCTCCT CATG

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 43 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GCATCGGGCC GCTAAGAATT TTCCTTTGTC ACCACATACA TAC

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 45 base pairs
  - (B) TYPE: nucleic acid

WO 93/10151

PCT/US92/10272

67

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GCATAGATCT GGGCTCAGAA TTTTCCTTG TCACCACATA CATAC

(2) INFORMATION FOR SEQ ID NO:17:

- (1) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 38 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GCATAGATCT GGGCTCTTCA ATTTCTCCTT GAGCAAC

WO 93/10151

PCT/US92/10272

68

CLAIMS

What is claimed is:

5

1. A receptor capable of binding oncostatin M and leukemia inhibitory factor, comprising gp130 covalently linked to LIF-R.


10

2. A receptor according to claim 1, wherein said receptor comprises a soluble gp130 polypeptide covalently linked to a soluble LIF-R polypeptide.

3. A receptor according to claim 1 wherein said receptor comprises gp130 covalently linked to LIF-R via a polypeptide linker.

15

4. A receptor according to claim 3, wherein said receptor is a recombinant fusion protein of the formula:



20

wherein  $R_1$  represents gp130;  $R_2$  represents LIF-R; and L represents a polypeptide linker.

25

5. A receptor according to claim 4 wherein the polypeptide linker comprises from 20 to 100 amino acids selected from the group consisting of glycine, asparagine, serine, threonine, and alanine.

30

6. A receptor according to claim 5 wherein the polypeptide linker comprises an amino acid sequence selected from the group consisting of:  $(Gly_4-Ser-Gly_5-Ser)_2$  and  $(Gly_4-Ser)_n$ , wherein n is 4-12.

7. An isolated DNA sequence encoding the receptor of claim 4.

8. A recombinant expression vector comprising the DNA sequence of claim 7.

35

9. A host cell containing the expression vector of claim 8.

WO 93/10151

PCT/US92/10272

69

10. A receptor according to claim 3, comprising a first fusion polypeptide that comprises an antibody Fc region polypeptide attached to the C-terminus of gp130, and a second fusion polypeptide that comprises an antibody Fc region polypeptide attached to the C-terminus of LIF-R, wherein said first fusion polypeptide is linked to said second fusion polypeptide *via* disulfide bonds between the Fc region polypeptides.

11. A receptor according to claim 1, 4, or 10, wherein:

a) said gp130 is encoded by an isolated DNA selected from the group consisting of a first DNA sequence comprising nucleotides 244-2369 of SEQ ID NO:1, a second DNA sequence comprising nucleotides 310-2369 of SEQ ID NO:1, and a third DNA sequence that will hybridize to said second DNA sequence under moderately stringent conditions; and

b) said LIF-R is encoded by an isolated DNA selected from the group consisting of a first DNA sequence comprising nucleotides 179-3182 of SEQ ID NO:5, a second DNA sequence comprising nucleotides 311-3182 of SEQ ID NO:5, and a third DNA sequence that will hybridize to said second DNA sequence under moderately stringent conditions.

12. A receptor according to claim 10 wherein said gp130 is a soluble gp130 polypeptide and said LIF-R is a soluble LIF-R polypeptide.

13. A fusion protein comprising an antibody Fc region polypeptide attached to the C-terminus of a soluble gp130 polypeptide.

14. An isolated DNA sequence encoding a fusion protein according to claim 13.

15. A fusion protein comprising an antibody Fc region polypeptide attached to the C-terminus of a soluble LIF-R polypeptide.

16. An isolated DNA sequence encoding a fusion protein according to claim 15.

17. A homodimeric receptor comprising two fusion proteins according to claim 15, linked *via* disulfide bonds between the Fc region polypeptides.

WO 93/10151

PCT/US92/10272

70

18. A process for preparing a receptor according to claim 4, comprising culturing a host cell transformed with an expression vector comprising a DNA sequence that encodes said fusion protein under conditions that promote expression of said fusion protein, and recovering said fusion protein.

19. A process for preparing a receptor according to claim 10, comprising culturing a host cell co-transfected with a first expression vector encoding said first fusion polypeptide and with a second expression vector encoding said second fusion polypeptide under conditions that promote expression of said first and second fusion polypeptides, and recovering said receptor.

20. A pharmaceutical composition for treating a disorder mediated by oncostatin M or LIF, comprising the receptor of claim 1, 4, 10, or 12, and a suitable diluent or carrier.



WO 93/10151


2/7

PCT/US92/10272

|   |             |                          |   |
|---|-------------|--------------------------|---|
| A | BIOG/LIFRGS | *OSM                     | □ |
| B | BIOG/LIFRGS | *OSM <sub>cell</sub> LIF | ■ |
| C | LIFRGS/cM   | *OSM                     | ○ |
| D | COS         | *OSM                     | ● |
| E | BIOG/cAV    | *OSM                     | △ |

 $K_D$  / sites per cellA.  $2.4 \times 10^{-10}$  M / 7,170B.  $5.5 \times 10^{-10}$  M / 1,150C/D.  $7.0 \times 10^{-11}$  M / 365E.  $5.0 \times 10^{-9}$  M / 3,408

FIGURE 2

 $\text{COS cells} / 10^5 \text{ LIF-OSM}$ 

WO 93/10151

3/7

PCT/US92/10272

FIGURE 3





FIGURE 4




WO 93/10151

PCT/US92/10272

5/7

FIGURE 5




WO 93/10151

PCT/US92/10272

6/7



FIGURE 6





**WO 93/10151**

PCT/L:S92/10272

7/7



## 7. ЗАКЛЮЧЕНИЕ



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US92/10272

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : C07K 13/00; C12N 15/62

US CL : 435/69.7; 514/2; 530/350

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/69.1, 69.7, 252.3, 320.1; 514/2; 530/350; 536/27

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, STN/MEDLINE

search terms: gp130, leukemia inhibitory factor, receptor

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages                                                                                                                                 | Relevant to claim No. |
|-----------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| A         | EMBO Journal, Vol. 10, No. 10, issued October 1991, D. P. Gearing et al, "Leukemia inhibitory factor receptor is structurally related to the IL-6 signal transducer, gp130", pages 2839-2848, see entire document. | 1-20                  |
| A         | Cell, Vol. 63, issued 21 December 1990, M. Hibi, et al, "Molecular Cloning and Expression of an IL-6 Signal Transducer, gp130", pages 1149-1157, see entire document.                                              | 1-20                  |
| A,P       | US, A, 5,155,027 (Sledziewski et al) 13 October 1992, column 5, line 38 to column 6, line 16.                                                                                                                      | 1-20                  |

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be part of particular relevance

\*P\* earlier documents published on or after the international filing date

\*L\* documents which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*E\* document published prior to the international filing date but later than the priority date claimed

\*T\*

later documents published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\*

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\*

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*A\*

document member of the same patent family

Date of the actual completion of the international search

05 March 1993

Date of mailing of the international search report

10 MAR 1993

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Authorized officer

JOHN D. ULM

Facsimile No. NOT APPLICABLE

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet) (July 1992)